



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/12, 5/10, C07K 14/705	A2	(11) International Publication Number: WO 99/43802 (43) International Publication Date: 2 September 1999 (02.09.99)
(21) International Application Number: PCT/JP99/00875 (22) International Filing Date: 25 February 1999 (25.02.99) (30) Priority Data: 10/46607 27 February 1998 (27.02.98) JP (71) Applicants (for all designated States except US): SAGAMI CHEMICAL RESEARCH CENTER [JP/JP]; 4-1, Nishi-Ohnuma 4-chome, Sagamihara-shi, Kanagawa 229-0012 (JP). PROTEGENE INC. [JP/JP]; 2-20-3, Naka-cho, Meguro-ku, Tokyo 153-0065 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): KATO, Seishi [JP/JP]; 3-46-50, Wakamatsu, Sagamihara-shi, Kanagawa 229-0014 (JP). SEKINE, Shingo [JP/JP]; Remonzu 101, 2-8-15, Atago, Ageo-shi, Saitama 362-0034 (JP). KIMURA, Tomoko [JP/JP]; 302, 4-1-28, Nishiikuta, Tama-ku, Kawasaki-shi, Kanagawa 214-0037 (JP). NAKAMURA, Nobuko [JP/JP]; 2-20-6, Kozonominami, Ayase-shi, Kanagawa 252-1122 (JP). (74) Agents: AOYAMA, Tamotsu et al.; Aoyama & Partners, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka 540-0001 (JP).		(81) Designated States: AU, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: HUMAN PROTEINS HAVING TRANSMEMBRANE DOMAINS AND DNAs ENCODING THESE PROTEINS (57) Abstract Human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic cells expressing said cDNAs. Said proteins and eucaryotic cells having said proteins on the membrane surface can be provided by expression of cDNAs coding for human proteins having transmembrane domains and of recombinants of these human cDNAs.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DESCRIPTION

Human Proteins Having Transmembrane
Domains and DNAs Encoding these Proteins

5

TECHINIAL FIELD

The present invention relates to human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic cells expressing said cDNAs. The proteins of the present invention can be employed as pharmaceuticals or as antigens for preparing antibodies against said proteins. The human cDNAs of the present invention can be utilized as probes for the gene diagnosis and gene sources for the gene therapy. Furthermore, the cDNAs can be utilized as gene sources for large-scale production of the proteins encoded by said cDNAs. Cells, wherein these membrane protein genes are introduced and membrane proteins are expressed in large amounts, can be utilized for detection of the corresponding ligands, screening of novel low-molecular pharmaceuticals, and so on.

20

BACKGROUND ART

Membrane proteins play important roles, as signal receptors, ion channels, transporters, etc. in the material transportation membrane and the information transmission which are mediated by the cell membrane. Examples thereof include receptors for a variety of cytokines, ion channels for the sodium ion, the potassium ion, the chloride ion, etc., transporters for saccharides and amino acids, and so on, where the genes of many of them have been cloned already.

30

It has been clarified that abnormalities of these membrane proteins are associated with a number of hitherto-cryptogenic diseases. For instance, a gene of a membrane protein having twelve transmembrane domains was identified as the gene responsible for cystic fibrosis [Rommens, J. M. et al., Science 245: 1059-1065 (1989)]. In addition, it has been clarified that several membrane proteins act as receptors when a virus infects the cells. For instance, HIV-1 is revealed to infect into the cells through mediation of a membrane protein fusin having a membrane protein on the T-cell membrane, a CD-4 antigen, and seven transmembrane domains [Feng, Y. et al., Science 272: 872-877 (1996)]. Therefore, discovery of a new membrane protein is anticipated to lead to elucidation of the causes of many diseases, so that isolation of a new gene coding for the membrane protein has been desired.

Heretofore, owing to difficulty in the purification, many membrane proteins have been isolated by an approach from the gene side. A general method is the so-called expression cloning which comprises transfection of a cDNA library in eucaryotic cells to express cDNAs and then detection of the cells expressing the target membrane protein on the membrane by an immunological technique using an antibody or a physiological technique on the change in the membrane permeability. However, this method is applicable only to cloning of a gene of a membrane protein with a known function.

In general, membrane proteins possess hydrophobic transmembrane domains inside the proteins, wherein, after synthesis thereof in the ribosome, these domains remain in the phospholipid membrane to be trapped in the membrane. Accordingly, the evidence of the cDNA for encoding the

membrane protein is provided by determination of the whole base sequence of a full-length cDNA followed by detection of highly hydrophobic transmembrane domains in the amino acid sequence of the protein encoded by said cDNA.

5

DISCLOSURE OF INVENTION

The object of the present invention is to provide novel human proteins having transmembrane domains, DNAs coding for said proteins, and expression vectors of said cDNAs as well as transformation eucaryotic cells that are capable of expressing said cDNAs.

As the result of intensive studies, the present inventors have been successful in cloning of cDNAs coding for proteins having transmembrane domains from the human full-length cDNA bank, thereby completing the present invention. In other words, the present invention provides human proteins having transmembrane domains, namely proteins containing any of the amino acid sequences represented by Sequence Nos. 1 to 7. Moreover, the present invention provides DNAs coding for the above-mentioned proteins, exemplified by cDNAs containing any of the base sequences represented by Sequence Nos. 8 to 15, 17, 19, 21, 23, 25 and 27, as well as transformation eucaryotic cells that are capable of expressing said cDNAs.

25

BRIEF DESCRIPTION OF DRAWINGS

Fig. 1: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP01434.

Fig. 2: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP01512.

Fig. 3: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP02080.

5 Fig. 4: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP02239.

Fig. 5: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP02375.

10 Fig. 6: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP10517.

15 Fig. 7: A figure depicting the hydrophobicity/hydrophilicity profile of the protein encoded by clone HP10521.

BEST MODE FOR CARRYING OUT THE INVENTION

The proteins of the present invention can be obtained, for example, by a method for isolation from human organs, cell lines, etc., a method for preparation of peptides by the chemical synthesis, or a method for production with the recombinant DNA technology using the DNAs coding for the transmembrane domains of the present invention, wherein the method for obtainment by the recombinant DNA technology is employed preferably. For instance, in vitro expression of the proteins can be achieved by preparation of an RNA by in vitro transcription from a vector having one of cDNAs of the present invention, followed by in vitro translation using this RNA as a template. Also, recombination of the translation region into a suitable expression vector by the method known in the art leads to expression of a large amount of the encoded protein by using prokaryotic cells

20

25

30

such as *Escherichia coli*, *Bacillus subtilis*, etc., and eucaryotic cells such as yeasts, insect cells, mammalian cells, etc.

5 In the case in which one of the proteins of the present invention is produced by expressing the DNA by in vitro translation, the protein of the present invention can be produced in vitro, when the translation region of said cDNA is subjected to recombination to a vector having an RNA polymerase promoter, followed by addition to an in vitro translation system such as a rabbit reticulocyte
10 lysate or a wheat germ extract, containing an RNA polymerase corresponding to the promoter. RNA polymerase inhibitors are exemplified by T7, T3, SP6, and the like. The vectors containing these RNA polymerase inhibitors are exemplified by pKA1, pCDM8, pT3/7 18, pT7/3 19, pBluescript
15 II, and so on. Furthermore, a membrane protein of the present invention can be expressed as the form incorporated in the microsome membrane, when a dog pancreas microsome or the like is added into the reaction system.

20 In the case in which a protein of the present invention is produced by expressing the DNA using a microorganism such as *Escherichia coli* etc., a recombinant expression vector bearing the translation region in the cDNA of the present invention is constructed in an expression vector having an origin, a promoter, a ribosome-binding site, a cDNA-cloning site, a terminator etc., which
25 can be replicated in the microorganism, and, after transformation of the host cells with said expression vector, the thus-obtained transformant is incubated, whereby the protein encoded by said cDNA can be produced on
30 a large scale in the microorganism. In this case, a protein fragment containing an optional region can be obtained by

carrying out the expression with inserting an initiation codon and a termination codon in front of and behind an optional translation region. Alternatively, a fusion protein with another protein can be expressed. Only a protein portion coding for said cDNA can be obtained by cleavage of said fusion protein with a suitable protease. The expression vector for *Escherichia coli* is exemplified by the pUC system, pBluescript II, the pET expression system, the pGEX expression system, and so on.

In the case in which one of the proteins of the present invention is produced by expressing the DNA in eucaryotic cells, the protein of the present invention can be produced as a transmembrane protein on the cell-membrane surface, when the translation region of said cDNA is subjected to recombination to an expression vector for eucaryotic cells that has a promoter, a splicing region, a poly(A) insertion site, etc., followed by introduction into the eucaryotic cells. The expression vector is exemplified by pKA1, pCDM8, pSVK3, pMSG, pSVL, pBK-CMV, pBK-RSV, EBV vector, pRS, pYES2, and so on. Examples of eucaryotic cells to be used in general include mammalian culture cells such as simian kidney cells COS7, Chinese hamster ovary cells CHO, etc., budding yeasts, fission yeasts, silkworm cells, *Xenopus laevis* egg cells, and so on, but any eucaryotic cells may be used, provided that they are capable of expressing the present proteins on the membrane surface. The expression vector can be introduced in the eucaryotic cells by methods known in the art such as the electroporation method, the potassium phosphate method, the liposome method, the DEAE-dextran method, and so on.

After one of the proteins of the present invention is expressed in prokaryotic cells or eucaryotic cells, the

objective protein can be isolated from the culture and purified by a combination of separation procedures known in the art. Such examples include treatment with a denaturing agent such as urea or a surface-active agent, sonication, enzymatic digestion, salting-out or solvent precipitation, dialysis, centrifugation, ultrafiltration, gel filtration, SDS-PAGE, isoelectric focusing, ion-exchange chromatography, hydrophobic chromatography, affinity chromatography, reverse phase chromatography, and so on.

The proteins of the present invention include peptide fragments (more than 5 amino acid residues) containing any partial amino acid sequence in the amino acid sequences represented by Sequence Nos. 1 to 7. These peptide fragments can be utilized as antigens for preparation of antibodies. Hereupon, among the proteins of the present invention, those having the signal sequence are secreted in the form of maturation proteins on the surface of the cells, after the signal sequences are removed. Therefore, these maturation proteins shall come within the scope of the present invention. The N-terminal amino acid sequences of the maturation proteins can be easily identified by using the method for the cleavage-site determination in a signal sequence [Japanese Patent Kokai Publication No. 1996-187100]. Furthermore, some membrane proteins undergo the processing on the cell surface to be converted to the secretory forms. Such proteins or peptides in the secretory forms shall come within the scope of the present invention. When sugar chain-binding sites are present in the amino acid sequences, expression in appropriate eucaryotic cells affords proteins wherein sugar chains are added. Accordingly, such proteins or peptides wherein sugar chains are added shall come within the scope of the present

invention.

The DNAs of the present invention include all DNAs coding for the above-mentioned proteins. Said DNAs can be obtained by using a method by chemical synthesis, a method
5 by cDNA cloning, and so on.

The cDNAs of the present invention can be cloned, for example, from cDNA libraries of the human cell origin. These cDNA are synthesized by using as templates poly(A)⁺ RNAs extracted from human cells. The human cells may be
10 cells delivered from the human body, for example, by the operation or may be the culture cells. The cDNAs can be synthesized by using any method selected from the Okayama-Berg method [Okayama, H. and Berg, P., Mol. Cell. Biol. 2: 161-170 (1982)], the Gubler-Hoffman method [Gubler, U. and
15 Hoffman, J. Gene 25: 263-269 (1983)], and so on, but it is preferred to use the capping method [Kato, S. et al., Gene 150: 243-250 (1994)], as exemplified in Examples, in order to obtain a full-length clone in an effective manner. In addition, commercially available, human cDNA libraries can
20 be utilized. Cloning of the cDNAs of the present invention from the cDNA libraries can be carried out by synthesis of an oligonucleotide on the basis of an optional portion in the cDNA base sequences of the present invention, followed by screening using this oligonucleotide as the probe
25 according to the colony or plaque hybridization by a method known in the art. In addition, the cDNA fragments of the present invention can be prepared by synthesis of an oligonucleotide to be hybridized at both termini of the objective cDNA fragment, followed by the usage of this
30 oligonucleotide as the primer for the RT-PCR method from an mRNA isolated from human cells.

The cDNAs of the present invention are characterized

by containing either of the base sequences represented by Sequence Nos. 8 to 14 or the base sequences represented by Sequence Nos. 15, 17, 19, 21, 23, 25 and 27. Table 1 summarizes the clone number (HP number), the cells
5 affording the cDNA, the total base number of the cDNA, and the number of the amino acid residues of the encoded protein, for each of the cDNAs.

Table 1

10

Sequence No.	HP No.	Cell	Number of bases	Number of amino acids
1, 8, 15	HP01434	Stomach Cancer	761	129
2, 9, 16	HP01512	Stomach Cancer	701	135
3, 10, 17	HP02080	Saos-2	393	79
4, 11, 18	HP02239	Stomach Cancer	1033	144
5, 12, 19	HP02375	PMA-U937	1270	282
6, 13, 20	HP10517	Liver	836	100
7, 14, 21	HP10521	Liver	1022	225

Hereupon, the same clones as the cDNAs of the present invention can be easily obtained by screening of the cDNA libraries constructed from the human cell lines and human
15 tissues utilized in the present invention by the use of an oligonucleotide probe synthesized on the basis of the cDNA base sequence described in any of Sequence Nos. 8 to 15, 17, 19, 21, 23, 25 and 27.

In general, the polymorphism due to the individual
20 difference is frequently observed in human genes. Accordingly, any cDNA that is subjected to insertion or deletion of one or plural nucleotides and/or substitution with other nucleotides in Sequence Nos. 8 to 15, 17, 19, 21, 23, 25 and 27 shall come within the scope of the present

invention.

In a similar manner, any protein that is formed by these modifications comprising insertion or deletion of one or plural amino acids and/or substitution with other amino acids shall come within the scope of the present invention, as far as the protein possesses the activity of any protein having the amino acid sequences represented by Sequence Nos. 1 to 7.

The cDNAs of the present invention include cDNA fragments (more than 10 bp) containing any partial base sequence in the base sequences represented by Sequence Nos. 8 to 14 or in the base sequences represented by Sequence Nos. 15, 17, 19, 21, 23, 25 and 27. Also, DNA fragments consisting of a sense chain and an anti-sense chain shall come within this scope. These DNA fragments can be utilized as the probes for the gene diagnosis.

In addition to the activities and uses described above, the polynucleotides and proteins of the present invention may exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified below. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or by administration or use of polynucleotides encoding such proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA).

Research Uses and Utilities

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein

is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on Southern gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The proteins provided by the present invention can similarly be used in assay to determine biological activity, including in a panel of multiple proteins for high-throughput screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the

corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands.

5 Where the protein binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these
10 binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for
15 commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory
20 Press, Sambrook, J., E.F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S.L. and A.R. Kimmel eds., 1987.

Nutritional Uses

25 Polynucleotides and proteins of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In
30 such cases the protein or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation,

such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the protein or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

5 Cytokine and Cell Proliferation/Differentiation Activity

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular

Immunology 133:327-341, 1991; Bertagnolli, et al., J. Immunol. 149:3778-3783, 1992; Bowman et al., J. Immunol. 152: 1756-1761, 1994.

5 Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and
10 Measurement of mouse and human Interferon γ , Schreiber, R.D. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of
15 hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley
20 and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6-Nordan, R. In Current Protocols in Immunology.
25 J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In Current Protocols in Immunology. J.E.e.a. Coligan
30 eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In Current

Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

Immune Stimulating or Suppressing Activity

A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the

present invention, including infections by HIV, hepatitis viruses, herpesviruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, a protein of the
5 present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example,
10 connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitis, myasthenia gravis, graft-versus-host disease and autoimmune
15 inflammatory eye disease. Such a protein of the present invention may also to be useful in the treatment of allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is
20 desired (including, for example, organ transplantation), may also be treatable using a protein of the present invention.

Using the proteins of the invention it may also be possible to immune responses, in a number of ways. Down
25 regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific
30 tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells

to the suppressive agent. Tolerance, which involves inducing non-responsiveness or anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as , for example, B7)), e.g., preventing high level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a molecule which inhibits or blocks interaction of a B7 lymphocyte antigen with its natural ligand(s) on immune cells (such as a soluble, monomeric form of a peptide having B7-2 activity alone or in conjunction with a monomeric form of a peptide having an activity of another B lymphocyte antigen (e.g., B7-1, B7-3) or blocking antibody), prior to transplantation can lead to the binding of the molecule to the natural ligand(s) on the immune cells without transmitting the corresponding costimulatory signal. Blocking B lymphocyte antigen function in this matter prevents cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover,

the lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular blocking reagents in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins in vivo as described in Lenschow et al., Science 257:789-792 (1992) and Turka et al., Proc. Natl. Acad. Sci USA, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of blocking B lymphocyte antigen function in vivo on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block costimulation of T cells by disrupting receptor:ligand interactions of B lymphocyte antigens can be used to

inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythmatosis in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., Fundamental Immunology, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (preferably a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response through stimulating B lymphocyte antigen function may be useful in cases of viral infection. In addition, systemic viral diseases such as influenza, the common cold, and encephalitis might be alleviated by the administration of stimulatory forms of B lymphocyte antigens systemically.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells in vitro with viral antigen-pulsed APCs either expressing a peptide of the present invention or together with a stimulatory form of a

soluble peptide of the present invention and reintroducing the in vitro activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells in vivo.

In another application, up regulation or enhancement of antigen function (preferably B lymphocyte antigen function) may be useful in the induction of tumor immunity. Tumor cells (e.g., sarcoma, melanoma, lymphoma, leukemia, neuroblastoma, carcinoma) transfected with a nucleic acid encoding at least one peptide of the present invention can be administered to a subject to overcome tumor-specific tolerance in the subject. If desired, the tumor cell can be transfected to express a combination of peptides. For example, tumor cells obtained from a patient can be transfected ex vivo with an expression vector directing the expression of a peptide having B7-2-like activity alone, or in conjunction with a peptide having B7-1-like activity and/or B7-3-like activity. The transfected tumor cells are returned to the patient to result in expression of the peptides on the surface of the transfected cell. Alternatively, gene therapy techniques can be used to target a tumor cell for transfection in vivo.

The presence of the peptide of the present invention having the activity of a B lymphocyte antigen(s) on the surface of the tumor cell provides the necessary costimulation signal to T cells to induce a T cell mediated

immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient amounts of MHC class I or MHC class II molecules, can be
5 transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I α chain protein and β_2 microglobulin protein or an MHC class II α chain protein and an MHC class II β chain protein to thereby express MHC class I or MHC class II
10 proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene
15 encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and
20 induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among
25 other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober,
30 Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans);

Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J.J. and Brunswick, M. In Current Protocols in Immunology. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551, 1991.

Hematopoiesis Regulating Activity

A protein of the present invention may be useful in

regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates
5 involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with
10 irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with
15 chemotherapy to prevent or treat consequent myelosuppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use
20 in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell
25 disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either in-vivo or ex-vivo (i.e.,
30 in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated

for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

5 Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

10 Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al., Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood 81:2903-2915, 1993.

15 Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., Experimental Hematology 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay,

Sutherland, H.J. In Culture of Hematopoietic Cells. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

Tissue Growth Activity

5 A protein of the present invention also may have utility in compositions used for bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions and ulcers.

10 A protein of the present invention, which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone fractures and cartilage damage or defects in humans and other animals. Such a preparation employing a protein of
15 the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced
20 craniofacial defects, and also is useful in cosmetic plastic surgery.

A protein of this invention may also be used in the treatment of periodontal disease, and in other tooth repair processes. Such agents may provide an environment to
25 attract bone-forming cells, stimulate growth of bone-forming cells or induce differentiation of progenitors of bone-forming cells. A protein of the invention may also be useful in the treatment of osteoporosis or osteoarthritis, such as through stimulation of bone and/or cartilage repair
30 or by blocking inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes.

Another category of tissue regeneration activity that may be attributable to the protein of the present invention is tendon/ligament formation. A protein of the present invention, which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues, and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide an environment to attract tendon or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors ex vivo for return in vivo to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The protein of the present invention may also be useful for proliferation of neural cells and for

regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a protein may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a protein of the invention.

Proteins of the invention may also be useful to promote better or faster closure of non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

It is expected that a protein of the present invention may also exhibit activity for generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A

protein of the invention may also exhibit angiogenic activity.

5 A protein of the present invention may also be useful for gut protection or regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

10 A protein of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

The activity of a protein of the invention may, among other means, be measured by the following methods:

15 Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

20 Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pps. 71-112 (Maibach, HI and Rovee, DT, eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

Activin/Inhibin Activity

25 A protein of the present invention may also exhibit activin- or inhibin-related activities. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins and are characterized by their ability to stimulate the release of
30 follicle stimulating hormone (FSH). Thus, a protein of the present invention, alone or in heterodimers with a member of the inhibin α family, may be useful as a contraceptive

based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the protein of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin- β group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, United States Patent 4,798,885. A protein of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as cows, sheep and pigs.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al., Nature 321:776-779, 1986; Mason et al., Nature 318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA 83:3091-3095, 1986.

Chemotactic/Chemokinetic Activity

A protein of the present invention may have chemotactic or chemokinetic activity (e.g., act as a chemokine) for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. Chemotactic and chemokinetic proteins can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic proteins provide particular advantages in treatment of wounds and

other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-146, 1995; Muller et al Eur. J. Immunol. 25: 1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153: 1762-1768, 1994.

Hemostatic and Thrombolytic Activity

A protein of the invention may also exhibit hemostatic or thrombolytic activity. As a result, such a protein is expected to be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A protein of the invention may also be useful for dissolving or inhibiting formation of thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke)).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis Res. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

Receptor/Ligand Activity

A protein of the present invention may also demonstrate activity as receptors, receptor ligands or inhibitors or agonists of receptor/ligand interactions. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of

cellular and humoral immune responses). Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in: Current Protocols in Immunology, Ed by J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1-7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160, 1989; Stoltenborg et al., J. Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

Anti-Inflammatory Activity

Proteins of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production of other factors which more directly inhibit or promote an inflammatory response. Proteins

exhibiting such activities can be used to treat inflammatory conditions including chronic or acute conditions), including without limitation inflammation associated with infection (such as septic shock, sepsis or
5 systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over
10 production of cytokines such as TNF or IL-1. Proteins of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material.

Tumor Inhibition Activity

In addition to the activities described above for
15 immunological treatment or prevention of tumors, a protein of the invention may exhibit other anti-tumor activities. A protein may inhibit tumor growth directly or indirectly (such as, for example, via ADCC). A protein may exhibit its tumor inhibitory activity by acting on tumor tissue or
20 tumor precursor tissue, by inhibiting formation of tissues necessary to support tumor growth (such as, for example, by inhibiting angiogenesis), by causing production of other factors, agents or cell types which inhibit tumor growth, or by suppressing, eliminating or inhibiting factors,
25 agents or cell types which promote tumor growth

Other Activities

A protein of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing,
30 infectious agents, including, without limitation, bacteria,

viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or
5 body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing,
10 utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive
15 disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in
20 the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and
25 the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

30 Examples

The present invention is embodied in more detail by

the following examples, but this embodiment is not intended to restrict the present invention. The basic operations and the enzyme reactions with regard to the DNA recombination are carried out according to the literature ["Molecular Cloning. A Laboratory Manual", Cold Spring Harbor Laboratory, 1989]. Unless otherwise stated, restrictive enzymes and a variety of modification enzymes to be used were those available from TAKARA SHUZO. The manufacturer's instructions were used for the buffer compositions as well as for the reaction conditions, in each of the enzyme reactions. The cDNA synthesis was carried out according to the literature [Kato, S. et al., Gene 150: 243-250 (1994)].

(1) Preparation of Poly(A)⁺ RNA

The histiocyte lymphoma cell line U937 (ATCC CRL 1593) stimulated by phorbol ester, the osteosarcoma cell line Saos-2 (ATCC HTB 85), tissues of stomach cancer delivered by the operation, and the liver were used for human cells to extract mRNAs. The cell line was incubated by a conventional procedure.

After about 1 g of the human cells was homogenized in 20 ml of a 5.5 M guanidinium thiocyanate solution, a total mRNA was prepared according to the literature [Okayama, H. et al., "Method in Enzymology", Vol. 164, Academic Press, 1987]. This was subjected to chromatography on oligo(dT)-cellulose column washed with a 20 mM Tris-hydrochloride buffer solution (pH 7.6), 0.5 M NaCl, and 1 mM EDTA to obtain a poly(A)⁺ RNA according to the above-described literature.

(2) Construction of cDNA Library

Ten micrograms of the above-mentioned poly(A)⁺ RNA were dissolved in a 100 mM Tris-hydrochloride buffer solution (pH 8), one unit of an RNase-free, bacterial

alkaline phosphatase was added, and the reaction was run at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 50 mM sodium acetate (pH 6), 1 mM EDTA, 0.1% 2-mercaptoethanol, and 0.01% Triton X-100. Thereto was added one unit of a tobacco-origin acid pyrophosphatase (Epicentre Technologies) and a total 100 µl volume of the resulting mixture was reacted at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in water to obtain a solution of a decapped poly(A)⁺ RNA.

The decapped poly(A)⁺ RNA and 3 nmol of a chimeric DNA-RNA oligonucleotide (5'-dG-dG-dG-dG-dA-dA-dT-dT-dC-dG-dA-G-G-A-3') were dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 7.5), 0.5 mM ATP, 5 mM MgCl₂, 10 mM 2-mercaptoethanol, and 25% polyethylene glycol, whereto was added 50 units of T4RNA ligase and a total 30 µl volume of the resulting mixture was reacted at 20°C for 12 hours. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in water to obtain a chimeric-oligo-capped poly(A)⁺ RNA.

After digestion of vector pKA1 (Japanese Patent Kokai Publication No. 1992-117292) developed by the present inventors with KpnI, about 60 dT tails were added using a terminal transferase. A vector primer to be used below was prepared by digestion of this product with EcoRV to remove a dT tail at one side.

After 6 µg of the previously-prepared chimeric-oligo-capped poly(A)⁺ RNA was annealed with 1.2 µg of the vector

primer, the resulting product was dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 8.3), 75 mM KCl, 3 mM MgCl₂, 10 mM dithiothreitol, and 1.25 mM dNTP (dATP + dCTP + dGTP + dTTP), 200 units of a reverse transcriptase (GIBCO-BRL) were added, and the reaction in a total 20 µl volume was run at 42°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 50 mM Tris-hydrochloride buffer solution (pH 7.5), 100 mM NaCl, 10 mM MgCl₂, and 1 mM dithiothreitol. Thereto were added 100 units of EcoRI and a total 20 µl volume of the resulting mixture was reacted at 37°C for one hour. After the reaction solution was subjected to phenol extraction, followed by ethanol precipitation, the resulting pellet was dissolved in a solution containing 20 mM Tris-hydrochloride buffer solution (pH 7.5), 100 mM KCl, 4 mM MgCl₂, 10 mM (NH₄)₂SO₄, and 50 µg/ml of the bovine serum albumin. Thereto were added 60 units of an *Escherichia coli* DNA ligase and the resulting mixture was reacted at 16°C for 16 hours. To the reaction solution were added 2 µl of 2 mM dNTP, 4 units of *Escherichia coli* DNA polymerase I, and 0.1 unit of *Escherichia coli* RNase H and the resulting mixture was reacted at 12°C for one hour and then at 22°C for one hour.

Next, the cDNA-synthesis reaction solution was used for transformation of *Escherichia coli* DH12S (GIBCO-BRL). The transformation was carried out by the electroporation method. A portion of the transformant was sprayed on the 2xYT agar culture medium containing 100 µg/ml ampicillin and the mixture was incubated at 37°C overnight. A colony formed on the agar medium was picked up at random and inoculated on 2 ml of the 2xYT culture medium containing

100 µg/ml ampicillin. After incubation at 37°C overnight, the culture mixture was centrifuged to separate the mycelia, from which a plasmid DNA was prepared by the alkaline lysis method. The plasmid DNA was subjected to double digestion
5 with EcoRI and NotI, followed by 0.8% agarose gel electrophoresis, to determine the size of the cDNA insert. Furthermore, using the thus-obtained plasmid as a template, the sequence reaction was carried out by using an M13
10 universal primer labeled with a fluorescent dye and a Taq polymerase (a kit of Applied Biosystems) and then the product was examined with a fluorescent DNA sequencer (Applied Biosystems) to determine an about 400-bp base sequence at the 5'-terminus of the cDNA. The sequence data were filed as the homo/protein cDNA bank database.

15 (3) Selection of cDNAs Encoding Proteins Having Transmembrane Domains

A base sequence registered in the homo/protein cDNA bank was converted to three frames of amino acid sequences and the presence or absence of an open reading frame (ORF)
20 beginning from the initiation codon was examined. Then, the selection was made for the presence of a signal sequence that is characteristic to a secretory protein at the N-terminus of the portion encoded by the ORF. These clones were sequenced from the both 5' and 3' directions by the
25 use of the deletion method using exonuclease III to determine the whole base sequence. The hydrophobicity/hydrophilicity profiles were obtained for proteins encoded by the ORF by the Kyte-Doolittle method [Kyte, J. & Doolittle, R. F., J. Mol. Biol. 157: 105-132
30 (1982)] to examine the presence or absence of a hydrophobic region. In the case in which there is a hydrophobic region of a putative transmembrane domain in the amino acid

sequence of an encoded protein, this protein was judged as a membrane protein.

(4) Functional Verification of Secretory Signal Sequence or Transmembrane Domains

5 It was verified by the method described in the literature [Yokoyama-Kobayashi, M. et al., Gene 163: 193-196 (1995)] that the N-terminal hydrophobic region in the secretory protein clone candidate obtained in the above-mentioned steps functions as a secretory signal sequence.
10 First, the plasmid containing the target cDNA was cleaved at an appropriate restriction enzyme site existing at the downstream of the portion expected for encoding the secretory signal sequence. In the case in which this restriction site was a protruding terminus, the site was
15 blunt-ended by the Klenow treatment or treatment with the T4DNA polymerase. Digestion with HindIII was further carried out and a DNA fragment containing the SV40 promoter and a cDNA encoding the secretory signal sequence at the downstream of the promoter was separated by agarose gel
20 electrophoresis. The resulting fragment was inserted between HindIII in pSSD3 (DDBJ/EMBL/GenBank Registration No. AB007632) and a restriction enzyme site selected so as to match with the urokinase-coding frame, thereby constructing a vector expressing a fusion protein of the secretory
25 signal sequence of the target cDNA and the urokinase protease domain.

After *Escherichia coli* (host: JM109) bearing the fusion-protein expression vector was incubated at 37°C for 2 hours in 2 ml of the 2xYT culture medium containing 100
30 µg/ml of ampicillin, the helper phage M13KO7 (50 µl) was added and the incubation was continued at 37°C overnight. A supernatant separated by centrifugation underwent

precipitation with polyethylene glycol to obtain single-stranded phage particles. These particles were suspended in 100 μ l of 1 mM Tris-0.1 mM EDTA, pH 8 (TE). Also, there were used as controls suspensions of single-stranded phage particles prepared in the same manner from pSSD3 and from the vector pKA1-UPA containing a full-length cDNA of urokinase [Yokoyama-Kobayashi, M. et al., Gene 163: 193-196 (1995)].

The culture cells originating from the simian kidney, COS7, were incubated at 37°C in the presence of 5% CO₂ in the Dulbecco's modified Eagle's culture medium (DMEM) containing 10% fetal calf albumin. Into a 6-well plate (Nunc Inc., 3 cm in the well diameter) were inoculated 1 X 10⁵ COS7 cells and incubation was carried out at 37°C for 22 hours in the presence of 5% CO₂. After the culture medium was removed, the cell surface was washed with a phosphate buffer solution and then washed again with DMEM containing 50 mM Tris-hydrochloric acid (pH 7.5) (TDMEM). To the resulting cells was added a suspension of 1 μ l of the single-stranded phage suspension, 0.6 ml of the DMEM culture medium, and 3 μ l of TRANSFECTAMTM (IBF Inc.) and the resulting mixture was incubated at 37°C for 3 hours in the presence of 5% CO₂. After the sample solution was removed, the cell surface was washed with TDMEM, 2 ml per well of DMEM containing 10% fetal calf albumin was added, and the incubation was carried out at 37°C for 2 days in the presence of 5% CO₂.

To 10 ml of 50 mM phosphate buffer solution (pH 7.4) containing 2% bovine fibrinogen (Miles Inc.), 0.5% agarose, and 1 mM calcium chloride were added 10 units of human thrombin (Mochida Pharmaceutical Co., Ltd.) and the resulting mixture was solidified in a plate of 9 cm in

diameter to prepare a fibrin plate. Ten microliters of the culture supernatant of the transfected COS7 cells were spotted on the fibrin plate, which was incubated at 37°C for 15 hours. In the case in which a clear circle appears on the fibrin plate, it is judged that the cDNA fragment codes for the amino acid sequence functioning as a secretory signal sequence. On the other hand, in case in which a clear circle is not formed, the cells were washed well, then the fibrin sheet was placed on the cells, and incubation was carried out at 37°C for 15 hours. In case in which a clear portion is formed on the fibrin sheet, it indicates that the urokinase activity was expressed on the cell surface. In other words, the cDNA fragment is judged to code for the transmembrane domains.

(5) Protein Synthesis by In Vitro Translation

The plasmid vector bearing the cDNA of the present invention was used for in vitro transcription/translation with a T_NT rabbit reticulocyte lysate kit (Promega). In this case, [³⁵S]methionine was added to label the expression product with a radioisotope. Each of the reactions was carried out according to the protocols attached to the kit. Two micrograms of the plasmid was reacted at 30°C for 90 minutes in a total 25 µl volume of the reaction solution containing 12.5 µl of T_NT rabbit reticulocyte lysate, 0.5 µl of a buffer solution (attached to kit), 2 µl of an amino acid mixture (methionine-free), 2 µl of [³⁵S]methionine (Amersham) (0.37 MBq/µl), 0.5 µl of T7RNA polymerase, and 20 U of RNasin. To 3 µl of the resulting reaction solution was added 2 µl of the SDS sampling buffer (125 mM Tris-hydrochloric acid buffer, pH 6.8, 120 mM 2-mercaptoethanol, 2% SDS solution, 0.025% bromophenol blue, and 20% glycerol) and the resulting

mixture was heated at 95°C for 3 minutes and then subjected to SDS-polyacrylamide gel electrophoresis. The molecular weight of the translation product was determined by carrying out the autoradiography.

5 (6) Expression by COS7

Escherichia coli bearing the expression vector of the protein of the present invention was infected with helper phage M13KO7 and single-stranded phage particles were obtained by the above-mentioned procedure. The thus-
10 obtained phage was used for introducing each expression vector in the culture cells originating from the simian kidney, COS7. After incubation at 37°C for 2 days in the presence of 5% CO₂, the incubation was continued for one hour in the culture medium containing [³⁵S]cystine or
15 [³⁵S]methionine. Collection and dissolution of the cells, followed by subjecting to SDS-PAGE, allowed to observe the presence of a band corresponding to the expression product of each protein, which did not exist in the COS7 cells.

(7) Clone Examples

20 <HP01434> (Sequence Nos. 1, 8, and 15)

Determination of the whole base sequence of the cDNA insert of clone HP01434 obtained from cDNA libraries of human stomach cancer revealed the structure consisting of a 69-bp 5'-nontranslation region, a 390-bp ORF, and a 302-bp
25 3'-nontranslation region. The ORF codes for a protein consisting of 129 amino acid residues and there existed one putative transmembrane domain. Figure 1 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro
30 translation resulted in formation of a translation product of 17 kDa that was a little larger than the molecular weight of 14,795 predicted from the ORF. Introduction of an

expression vector, wherein the HindIII-PstI (blunt-ended by treatment with T4RNA polymerase) fragment containing a cDNA portion coding for the N-terminal 68 amino acid residues of the present protein was inserted into the HindIII-PmaCI site of pSSD3, into the COS7 cells revealed that the urokinase activity was not detectable in the culture medium to indicate that the present protein remains in the membrane.

The search of the protein data base by using the amino acid sequence of the present protein revealed that the protein was analogous to the murine FK506-binding protein (SWISS-PROT Accession No. P45878). Table 2 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the murine FK506-binding protein (MM). Therein, the marks of -, *, and . represent a gap, an amino acid residue identical with the protein of the present invention, and an amino acid residue analogous to the protein of the present invention, respectively. The both proteins possessed a homology of 46.8% in the 94 amino acid residues.

Table 2

25	HS MHFLFRFIVFFYLWGLFTAQRQKKEESTEEVKIEVLHRPENCSKTSKKGDLLNAHYDGYL
 * ..* .*.***.*. **.* *
	MM MRLSWILTILSICLSALAAATGAEGKRKLQIGVKRVDHCPIKSRKGDVLHMHYTGKL
	HS AKDGSKFYCSRTQNEGHPKWFVLGVGQVIKGLDIAMTDMCPGEKRKVVIPPSFAYGKEGY
	.***.* . * .** . * * **.****** * .. .** *****.***.....* *
	MM -EDGTEFDSSLPQN--QPFVFSLGTQGVIKGWDQGLLGMCEGEKRKLVIPSELGYGERGA
30	HS DKPLLAKGI
	MM PPKIPGGATLVFEVELLKIERRSEL

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the registration of sequences that possessed a homology of 90% or more (for example, Accession No. AA431230) in EST, but any of the
5 sequences was shorter than the present cDNAs and was not found to contain the initiation codon.

The murine FK506-binding protein possesses the peptidyl prolysis-transisomerase activity and is associated with the folding of proteins [Hendrickson, B. A.
10 et al., Gene 134: 271-275 (1993)].

<HP01512> (Sequence Nos. 2, 9, and 17)

Determination of the whole base sequence of the cDNA insert of clone HP01512 obtained from cDNA libraries of human stomach cancer revealed the structure consisting of a
15 45-bp 5'-nontranslation region, a 408-bp ORF, and a 248-bp 3'-nontranslation region. The ORF codes for a protein consisting of 135 amino acid residues and there existed one putative transmembrane domain. Figure 2 depicts the hydrophobicity/hydrophilicity profile, obtained by the
20 Kyte-Doolittle method, of the present protein.

The search of the protein data base by using the amino acid sequence of the present protein revealed that the protein was analogous to the nematode imaginary protein W02B12.7 (PID Accession No. 1044857). Table 3 shows the
25 comparison of the amino acid sequence between the human protein of the present invention (HP) and the nematode imaginary protein W02B12.7 (CE). Therein, the marks of -, *, and . represent a gap, an amino acid residue identical with the protein of the present invention, and an amino acid
30 residue analogous to the protein of the present invention, respectively. The C-terminal 87 amino acid residues possessed a homology of 49.4% with the N-terminal side of

the nematode imaginary protein W02B12.7.

Table 3

5	HS	MVLESVARIVKVQLPAYLKRLPVPEITGTFARLTVSEWLRLLPFLGVLALLGYLA
		*. **.****.
	CE	MTIAGFCALSLIQDHCCWTTTPQHCSVAELGTMPCPTQVSGRCVATTA AVLAGGALIGYLV
	HS	VRPFLPKKKQOKDSLINLKIQKENPKVVNEINIEDLCLTKAAYCRCWRSKTFPACDGSNH
		* *... * *** .. *.****.****. *.****.****.****.
10	CE	GYKF-----GQRSARCNYKIQLDNKNIVDTVDIEDIG-EKKAFCRCWKSEKWPYCDGSHG
	HS	KHNELTGDNVGPLILKKKEV
		. *.****.*
	CE	KHNKETGDNVGPLIVKSEKNLYIYIIISDFYNNHTNDLKHQIAQLERKTATIPKLENQLH

15

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. AA429420) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

20

<HP02080> (Sequence Nos. 3, 10, and 19)

25

Determination of the whole base sequence of the cDNA insert of clone HP02080 obtained from cDNA libraries of human osteosarcoma cell line Saos-2 revealed the structure consisting of an 80-bp 5'-nontranslation region, a 240-bp ORF, and a 73-bp 3'-nontranslation region. The ORF codes for a protein consisting of 79 amino acid residues and there existed one putative transmembrane domain. Figure 3 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation

30

product of 10 kDa that was almost identical with the molecular weight of 8,177 predicted from the ORF.

The search of the protein data base using the amino acid sequence of the present protein has revealed that the protein was analogous to a yeast imaginary protein Lpg10p (PID Accession No. 1749572). Table 4 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the yeast imaginary protein Lpg10p (SC). Therein, the marks of -, *, and . represent a gap, an amino acid residue identical with the protein of the present invention, and an amino acid residue analogous to the protein of the present invention, respectively. The both proteins possessed a homology of 46.3% in the entire region.

Table 4

	HS	MPVAVGPYQSQPSCFDRVKMGFVMGCAVGMAGALFGTFSCLRIGMRGRELMGGIGKTM
		*** *..*** .**.*.*.. * ***. . ** * *... ..* *
20	SC	MQSMQPSTVDKLMGAIMGSAAGLGIGFLFGGVAVLRYGPGPRGFLRTLQOYM
	HS	MQSGGTFGTFMAIGMGIRC
		. *..*** **.* **
	SC	L TSAATFGFFMSIGSVIRNEDIPLIQSGSHWNQRLLENANSSRIFALAMQOAKSSPRK

25

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the registration of sequences that possessed a homology of 90% or more (for example, Accession No. AA348987) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

30

<HP02239> (Sequence Nos. 4, 11, and 21)

Determination of the whole base sequence of the cDNA insert of clone HP02239 obtained from cDNA libraries of human stomach cancer revealed the structure consisting of a
5 47-bp 5'-nontranslation region, a 435-bp ORF, and a 551-bp
3'-nontranslation region. The ORF codes for a protein consisting of 144 amino acid residues and there existed three putative transmembrane domains. Figure 4 depicts the hydrophobicity/hydrophilicity profile, obtained by the
10 Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product of 17 kDa that was almost identical with the molecular weight of 16,687 predicted from the ORF.

The search of the protein data base using the amino
15 acid sequence of the present protein has revealed that the protein was analogous to mouse cornichon (PID Accession No. 2460430). Table 5 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the mouse cornichon (MM). Therein, the marks of *
20 and . represent an amino acid residue identical with the protein of the present invention and an amino acid residue analogous to the protein of the present invention, respectively. The both proteins possessed a homology of 99.3% in the entire region.

Table 5

	HS	MAFTFAAFCYMLALLLTAALIFFAIWHIIAFDELKTDYKNPIDQCNTLNPLVLPEYLIHA

5	MM	MAFTFAAFCYMLALLLTAALIFFAIWHIIAFDELKTDYKNPIDQCNTLNPLVLPEYLIHA
	HS	FFCVMFLCAAEWLTGLNMPLLAYHIWRYMSRPVMSGPGLYDPTTIMNADILAYCQKEGW
		*****.*****
	MM	FFCVMFLCAAEWLTGLNMPLLAYHIWRYMSRPVSAPGLYDPTTIMNADILAYCQKEGW
	HS	CKLAFYLLAFFYYLYGMIYVLVSS
10		*****
	MM	CKLAFYLLAFFYYLYGMIYVLVSS

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. W02973) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

Cornichon has been found in the genesis of *Drosophila* as a membrane protein that is essential to the morphogenesis [Roth, S. et al., Cell 81:967-978 (1995)].

<HP02375> (Sequence Nos. 5, 12, and 23)

Determination of the whole base sequence of the cDNA insert of clone HP02375 obtained from cDNA libraries of human lymphoma cell line U937 revealed the structure consisting of an 85-bp 5'-nontranslation region, an 849-bp ORF, and a 336-bp 3'-nontranslation region. The ORF codes for a protein consisting of 282 amino acid residues and possessed a signal-like sequence at the N-terminus and one transmembrane domain at the C-terminus. Accordingly, the present protein is considered to be a type-I membrane

protein. Figure 5 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein.

5 The search of the protein data base using the amino acid sequence of the present protein has revealed that this had an analogy to the human LDL-related protein 1 (PID Accession No. 1708865). Table 6 shows the comparison of the amino acid sequence between the human protein of the present invention (HP) and the human LDL-related protein 1
10 (LR). Therein, the marks of -, *, and . represent a gap, an amino acid residue identical with the protein of the present invention, and an amino acid residue analogous to the protein of the present invention, respectively. An intermediate portion of the present protein possessed a
15 homology of 36.4% with the LDL receptor class-A domain portion of the human LDL-related protein 1. Particularly, the position of cysteine was preserved.

Table 6

```

HS                                     MSGGWMAQVGAW

5   LR CGDRSDESASCAYPTCFPLTQFTCNNGRCININWRCNDNDNDCGNSDEAGCSHSCSSSTQF
    HS RTGALGLALLLLGLGLEAAASPLSTPSTAQAAGPSSGSCPPTKFCRTSGLCVPLTW
        .....*.~.*.~.....**** .***.~~ *
    LR KCNSGRCIPEHWTCGDNDNCGDYSETHANCTNQATRPPGGCHTDEFQCRLDGLCIPLRW
    HS RCDRDLDCSDGSDEEECR--IEPCTQKGQ--C-PPPPGLPCP--CTGVSDCSGGTDKKLR
10   *** ~ ** *.***..* .. *... . * ... .. *.~ **.~....*..
    LR RCDGDTDCMDSSDEKSCEGVTHVCDPSVKFGCKDSARCISKAWCDGDNDCEDNSDEE--
    HS NCSR LACLAGE LRCTL SDD-CIPLTW RCDGH PDCPDSSDELGC GTNEILPEG DATIMGPP
        **..*** . . *. ... *.~ . ***. ~~ *.***
    LR NCESLACRP PSHPCANN TSVCLPPDKLCDGNDDCGDGSDEGELCDQCSLNNGGCSHNCSV
15   HS VTLESVTSLRNATTMGP PVTL ESVP SVGNATSSSAGD QSGSP TAYGVIAAAAVLSASLVT

    LR APGEGIVC SCPLGMELGPDNH TCQIQSYCAKHLKCSQKCDQN KFSVKCSCYEGWVLEPDG
    HS ATLLLLSWLRAQERLRPLGLLVAMKESLLLSEQKTSLP

20   LR ESCRSILD PFKPF IIFS NRHEIRRIDLHKGDYSVLVPGLRNTIALDFHLSQSALYWTDVVE

```

Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the registration of sequences that possessed a homology of 90% or more (for example, Accession No. H94922) in EST, but any of them was shorter than the present cDNA and was found not to contain the initiation codon.

30 The human LDL-related protein 1 has been shown to be an α 2-macroglobulin receptor [Kristensen, T. et al., FEBS Lett. 276: 151-155 (1990)]. The protein possessing such an LDL receptor class-A domain is considered to work as a receptor for proteins existing in the serum.

<HP10517> (Sequence Nos. 6, 13, and 25)

Determination of the whole base sequence of the cDNA insert of clone HP10517 obtained from cDNA libraries of human liver revealed the structure consisting of a 163-bp 5'-nontranslation region, a 303-bp ORF, and a 370-bp 3'-nontranslation region. The ORF codes for a protein consisting of 100 amino acid residues and possessed one transmembrane domain. Figure 6 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product of 12 kDa that was almost identical with the molecular weight of 11,796 predicted from the ORF.

The search of the protein data base using the amino acid sequence of the present protein has not identified any known protein having an analogy. Furthermore, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. R68523) in EST, but, since they are partial sequences, it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

<HP10521> (Sequence Nos. 7, 14, and 27)

Determination of the whole base sequence of the cDNA insert of clone HP10521 obtained from cDNA libraries of human liver revealed the structure consisting of a 55-bp 5'-nontranslation region, a 678-bp ORF, and a 289-bp 3'-nontranslation region. The ORF codes for a protein consisting of 225 amino acid residues and possessed four transmembrane domains. Figure 7 depicts the hydrophobicity/hydrophilicity profile, obtained by the Kyte-Doolittle method, of the present protein. In vitro translation resulted in formation of a translation product

of 27 kDa that was a little larger than the molecular weight of 24,809 predicted from the ORF.

5 The search of the protein data base using the amino acid sequence of the present protein has not revealed the presence of any known protein having an analogy. Also, the search of the GenBank using the base sequences of the present cDNA has revealed the presence of sequences that possessed a homology of 90% or more (for example, Accession No. AA043627) in EST, but, since they are partial sequences,
10 it can not be judged whether or not any of these sequences codes for the same protein as the protein of the present invention.

INDUSTRIAL APPLICABILITY

15 The present invention provides human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic cells expressing said cDNAs. All of the proteins of the present invention exist in the cell membrane, so that they
20 are considered to be proteins controlling the proliferation and the differentiation of the cells. Accordingly, the proteins of the present invention can be employed as pharmaceuticals such as carcinostatic agents relating to the control of the proliferation and the differentiation of
25 the cells or as antigens for preparing antibodies against said proteins. The cDNAs of the present invention can be utilized as probes for the gene diagnosis and gene sources for the gene therapy. Furthermore, the cDNAs can be utilized for large-scale expression of said proteins. Cells,
30 wherein these membrane protein genes are introduced to possess said proteins on the membrane surface, can be utilized for detection of the corresponding ligands,

screening of novel low-molecular pharmaceuticals, and so on.

The present invention also provides genes corresponding to the polynucleotide sequences disclosed herein. "Corresponding genes" are the regions of the genome that are transcribed to produce the mRNAs from which cDNA polynucleotide sequences are derived and may include contiguous regions of the genome necessary for the regulated expression of such genes. Corresponding genes may therefore include but are not limited to coding sequences, 5' and 3' untranslated regions, alternatively spliced exons, introns, promoters, enhancers, and silencer or suppressor elements. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. An "isolated gene" is a gene that has been separated from the adjacent coding sequences, if any, present in the genome of the organism from which the gene was isolated.

Organisms that have enhanced, reduced, or modified expression of the gene(s) corresponding to the polynucleotide sequences disclosed herein are provided. The desired change in gene expression can be achieved through the use of antisense polynucleotides or ribozymes that bind and/or cleave the mRNA transcribed from the gene (Albert and Morris, 1994, Trends Pharmacol. Sci. 15(7): 250-254; Lavarosky et al., 1997, Biochem. Mol. Med. 62(1): 11-22; and Hampel, 1998, Prog. Nucleic Acid Res. Mol. Biol. 58: 1-39; all of which are incorporated by reference herein). Transgenic animals that have multiple copies of

the gene(s) corresponding to the polynucleotide sequences disclosed herein, preferably produced by transformation of cells with genetic constructs that are stably maintained within the transformed cells and their progeny, are provided. Transgenic animals that have modified genetic control regions that increase or reduce gene expression levels, or that change temporal or spatial patterns of gene expression, are also provided (see European Patent No. 0 649 464 B1, incorporated by reference herein). In addition, organisms are provided in which the gene(s) corresponding to the polynucleotide sequences disclosed herein have been partially or completely inactivated, through insertion of extraneous sequences into the corresponding gene(s) or through deletion of all or part of the corresponding gene(s). Partial or complete gene inactivation can be accomplished through insertion, preferably followed by imprecise excision, of transposable elements (Plasterk, 1992, Bioessays 14(9): 629-633; Zwaal et al., 1993, Proc. Natl. Acad. Sci. USA 90(16): 7431-7435; Clark et al., 1994, Proc. Natl. Acad. Sci. USA 91(2): 719-722; all of which are incorporated by reference herein), or through homologous recombination, preferably detected by positive/negative genetic selection strategies (Mansour et al., 1988, Nature 336: 348-352; U.S. Patent Nos. 5,464,764; 5,487,992; 5,627,059; 5,631,153; 5,614,396; 5,616,491; and 5,679,523; all of which are incorporated by reference herein). These organisms with altered gene expression are preferably eukaryotes and more preferably are mammals. Such organisms are useful for the development of non-human models for the study of disorders involving the corresponding gene(s), and for the development of assay systems for the identification of molecules that interact with the protein product(s) of

the corresponding gene(s).

Where the protein of the present invention is membrane-bound (e.g., is a receptor), the present invention also provides for soluble forms of such protein. In such
5 forms part or all of the intracellular and transmembrane domains of the protein are deleted such that the protein is fully secreted from the cell in which it is expressed. The intracellular and transmembrane domains of proteins of the invention can be identified in accordance with known
10 techniques for determination of such domains from sequence information.

Proteins and protein fragments of the present invention include proteins with amino acid sequence lengths that are at least 25% (more preferably at least 50%, and
15 most preferably at least 75%) of the length of a disclosed protein and have at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with that disclosed protein, where sequence identity is determined by comparing the amino acid
20 sequences of the proteins when aligned so as to maximize overlap and identity while minimizing sequence gaps. Also included in the present invention are proteins and protein fragments that contain a segment preferably comprising 8 or more (more preferably 20 or more, most preferably 30 or
25 more) contiguous amino acids that shares at least 75% sequence identity (more preferably, at least 85% identity; most preferably at least 95% identity) with any such segment of any of the disclosed proteins.

Species homologs of the disclosed polynucleotides and
30 proteins are also provided by the present invention. As used herein, a "species homologue" is a protein or polynucleotide with a different species of origin from that

of a given protein or polynucleotide, but with significant sequence similarity to the given protein or polynucleotide, as determined by those of skill in the art. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species.

The invention also encompasses allelic variants of the disclosed polynucleotides or proteins; that is, naturally-occurring alternative forms of the isolated polynucleotide which also encode proteins which are identical, homologous, or related to that encoded by the polynucleotides.

The invention also includes polynucleotides with sequences complementary to those of the polynucleotides disclosed herein.

The present invention also includes polynucleotides capable of hybridizing under reduced stringency conditions, more preferably stringent conditions, and most preferably highly stringent conditions, to polynucleotides described herein. Examples of stringency conditions are shown in the table below: highly stringent conditions are those that are at least as stringent as, for example, conditions A-F; stringent conditions are at least as stringent as, for example, conditions G-L; and reduced stringency conditions are at least as stringent as, for example, conditions M-R.

Table 7

Stringency Condition	Polynucleotide Hybrid	Hybrid Length (bp) [‡]	Hybridization Temperature and Buffer [†]	Wash Temperature and Buffer [†]
A	DNA : DNA	≥50	65°C; 1×SSC -or- 42°C; 1×SSC, 50% formamide	65°C; 0.3×SSC
B	DNA : DNA	<50	T _B *; 1×SSC	T _B *; 1×SSC
C	DNA : RNA	≥50	67°C; 1×SSC -or- 45°C; 1×SSC, 50% formamide	67°C; 0.3×SSC
D	DNA : RNA	<50	T _D *; 1×SSC	T _D *; 1×SSC
E	RNA : RNA	≥50	70°C; 1×SSC -or- 50°C; 1×SSC, 50% formamide	70°C; 0.3×SSC
F	RNA : RNA	<50	T _F *; 1×SSC	T _F *; 1×SSC
G	DNA : DNA	≥50	65°C; 4×SSC -or- 42°C; 4×SSC, 50% formamide	65°C; 1×SSC
H	DNA : DNA	<50	T _H *; 4×SSC	T _H *; 4×SSC
I	DNA : RNA	≥50	67°C; 4×SSC -or- 45°C; 4×SSC, 50% formamide	67°C; 1×SSC
J	DNA : RNA	<50	T _J *; 4×SSC	T _J *; 4×SSC
K	RNA : RNA	≥50	70°C; 4×SSC -or- 50°C; 4×SSC, 50% formamide	67°C; 1×SSC
L	RNA : RNA	<50	T _L *; 2×SSC	T _L *; 2×SSC
M	DNA : DNA	≥50	50°C; 4×SSC -or- 40°C; 6×SSC, 50% formamide	50°C; 2×SSC
N	DNA : DNA	<50	T _N *; 6×SSC	T _N *; 6×SSC
O	DNA : RNA	≥50	55°C; 4×SSC -or- 42°C; 6×SSC, 50% formamide	55°C; 2×SSC
P	DNA : RNA	<50	T _P *; 6×SSC	T _P *; 6×SSC
Q	RNA : RNA	≥50	60°C; 4×SSC -or- 45°C; 6×SSC, 50% formamide	60°C; 2×SSC
R	RNA : RNA	<50	T _R *; 4×SSC	T _R *; 4×SSC

5 ‡ : The hybrid length is that anticipated for the hybridized region(s) of the hybridizing polynucleotides. When hybridizing a polynucleotide to a target polynucleotide of unknown sequence, the hybrid length is assumed to be that of the hybridizing polynucleotide. When polynucleotides of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the polynucleotides and identifying the region or regions of optimal sequence complementarity.

10 † : SSPE (1×SSPE is 0.15M NaCl, 10mM NaH₂PO₄, and 1.25mM EDTA, pH7.4) can be substituted for SSC (1×SSC is 0.15M NaCl and 15mM sodium citrate) in the hybridization and wash buffers; washes are performed for 15 minutes after hybridization is complete.

*T_B - T_R : The hybridization temperature for hybrids anticipated to be less than 50 base

pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, $T_m(^{\circ}\text{C})=2(\text{\# of A + T bases}) + 4(\text{\# of G + C bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}\text{C})=81.5 + 16.6(\log_{10}[\text{Na}^+]) + 0.41 (\% \text{G+C}) - (600/N)$, where N is the number of bases in the hybrid, and $[\text{Na}^+]$ is the concentration of sodium ions in the hybridization buffer ($[\text{Na}^+]$ for $1\times\text{SSC}=0.165\text{M}$).

Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and Current Protocols in Molecular Biology, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Preferably, each such hybridizing polynucleotide has a length that is at least 25% (more preferably at least 50%, and most preferably at least 75%) of the length of the polynucleotide of the present invention to which it hybridizes, and has at least 60% sequence identity (more preferably, at least 75% identity; most preferably at least 90% or 95% identity) with the polynucleotide of the present invention to which it hybridizes, where sequence identity is determined by comparing the sequences of the hybridizing polynucleotides when aligned so as to maximize overlap and identity while minimizing sequence gaps.

CLAIMS

1. Proteins containing any of the amino acid sequences represented by Sequence Nos. 1 to 7.
- 5 2. DNAs coding for any of the proteins described in Claim 1.
3. cDNAs containing any of the base sequences represented by Sequence Nos. 8 to 14.
4. The cDNAs described in Claim 3 comprising any of
10 the base sequences represented by Sequence Nos. 15, 17, 19, 21, 23, 25 and 27.
5. Expression vectors that are capable of expressing any of the DNAs described in Claim 2 to Claim 4 by in vitro translation or in eucaryotic cells
- 15 6. Transformation eucaryotic cells that are capable of expressing any of the DNAs described in Claim 2 to Claim 4 and of producing the proteins described in Claim 1.

1/7




Fig. 1

2/7




Fig. 2

3/7




Fig. 3

4/7




Fig. 4




Fig. 5

6/7




Fig. 6

7/7




Fig. 7

1/27

SEQUENCE LISTING

<110> Sagami Chemical Research Center et al.

5 <120> Human Proteins Having Transmembrane Domains and DNAs Encoding these
Proteins

<130> 661098

10 <141> 1999-02-25

<150> JP 10-046607

<151> 1998-02-27

15 <160> 28

<170> Windows 95 (Word 98)

<210> 1

20 <211> 129

<212> PRT

<213> Homo sapiens

<400> 1

25 Met His Phe Leu Phe Arg Phe Ile Val Phe Phe Tyr Leu Trp Gly Leu

1 5 10 15

Phe Thr Ala Gln Arg Gln Lys Lys Glu Glu Ser Thr Glu Glu Val Lys

20 25 30

Ile Glu Val Leu His Arg Pro Glu Asn Cys Ser Lys Thr Ser Lys Lys

30 35 40 45

Gly Asp Leu Leu Asn Ala His Tyr Asp Gly Tyr Leu Ala Lys Asp Gly

50 55 60

2/27

Ser Lys Phe Tyr Cys Ser Arg Thr Gln Asn Glu Gly His Pro Lys Trp
 65 70 75 80
 Phe Val Leu Gly Val Gly Gln Val Ile Lys Gly Leu Asp Ile Ala Met
 85 90 95
 5 Thr Asp Met Cys Pro Gly Glu Lys Arg Lys Val Val Ile Pro Pro Ser
 100 105 110
 Phe Ala Tyr Gly Lys Glu Gly Tyr Asp Lys Pro Leu Leu Ala Lys Gly
 115 120 125
 Ile
 10
 <210> 2
 <211> 135
 <212> PRT
 <213> Homo sapiens
 15
 <400> 2
 Met Val Leu Glu Ser Val Ala Arg Ile Val Lys Val Gln Leu Pro Ala
 1 5 10 15
 Tyr Leu Lys Arg Leu Pro Val Pro Glu Ser Ile Thr Gly Phe Ala Arg
 20 20 25 30
 Leu Thr Val Ser Glu Trp Leu Arg Leu Leu Pro Phe Leu Gly Val Leu
 35 40 45
 Ala Leu Leu Gly Tyr Leu Ala Val Arg Pro Phe Leu Pro Lys Lys Lys
 50 55 60
 25 Gln Gln Lys Asp Ser Leu Ile Asn Leu Lys Ile Gln Lys Glu Asn Pro
 65 70 75 80
 Lys Val Val Asn Glu Ile Asn Ile Glu Asp Leu Cys Leu Thr Lys Ala
 85 90 95
 Ala Tyr Cys Arg Cys Trp Arg Ser Lys Thr Phe Pro Ala Cys Asp Gly
 100 105 110
 30 Ser His Asn Lys His Asn Glu Leu Thr Gly Asp Asn Val Gly Pro Leu
 115 120 125

3/27

Ile Leu Lys Lys Lys Glu Val

130

135

<210> 3

5

<211> 79

<212> PRT

<213> Homo sapiens

<400> 3

10

Met Pro Val Ala Val Gly Pro Tyr Gly Gln Ser Gln Pro Ser Cys Phe

1

5

10

15

Asp Arg Val Lys Met Gly Phe Val Met Gly Cys Ala Val Gly Met Ala

20

25

30

Ala Gly Ala Leu Phe Gly Thr Phe Ser Cys Leu Arg Ile Gly Met Arg

15

35

40

45

Gly Arg Glu Leu Met Gly Gly Ile Gly Lys Thr Met Met Gln Ser Gly

50

55

60

Gly Thr Phe Gly Thr Phe Met Ala Ile Gly Met Gly Ile Arg Cys

65

70

75

20

<210> 4

<211> 144

<212> PRT

<213> Homo sapiens

25

<400> 4

Met Ala Phe Thr Phe Ala Ala Phe Cys Tyr Met Leu Ala Leu Leu Leu

1

5

10

15

Thr Ala Ala Leu Ile Phe Phe Ala Ile Trp His Ile Ile Ala Phe Asp

30

20

25

30

Glu Leu Lys Thr Asp Tyr Lys Asn Pro Ile Asp Gln Cys Asn Thr Leu

35

40

45

4/27

Asn Pro Leu Val Leu Pro Glu Tyr Leu Ile His Ala Phe Phe Cys Val
 50 55 60
 Met Phe Leu Cys Ala Ala Glu Trp Leu Thr Leu Gly Leu Asn Met Pro
 65 70 75 80
 5 Leu Leu Ala Tyr His Ile Trp Arg Tyr Met Ser Arg Pro Val Met Ser
 85 90 95
 Gly Pro Gly Leu Tyr Asp Pro Thr Thr Ile Met Asn Ala Asp Ile Leu
 100 105 110
 Ala Tyr Cys Gln Lys Glu Gly Trp Cys Lys Leu Ala Phe Tyr Leu Leu
 10 115 120 125
 Ala Phe Phe Tyr Tyr Leu Tyr Gly Met Ile Tyr Val Leu Val Ser Ser
 130 135 140

 <210> 5
 15 <211> 282
 <212> PRT
 <213> Homo sapiens

 <400> 5
 20 Met Ser Gly Gly Trp Met Ala Gln Val Gly Ala Trp Arg Thr Gly Ala
 1 5 10 15
 Leu Gly Leu Ala Leu Leu Leu Leu Leu Gly Leu Gly Leu Gly Leu Glu
 20 25 30
 Ala Ala Ala Ser Pro Leu Ser Thr Pro Thr Ser Ala Gln Ala Ala Gly
 25 35 40 45
 Pro Ser Ser Gly Ser Cys Pro Pro Thr Lys Phe Gln Cys Arg Thr Ser
 50 55 60
 Gly Leu Cys Val Pro Leu Thr Trp Arg Cys Asp Arg Asp Leu Asp Cys
 65 70 75 80
 30 Ser Asp Gly Ser Asp Glu Glu Glu Cys Arg Ile Glu Pro Cys Thr Gln
 85 90 95
 Lys Gly Gln Cys Pro Pro Pro Pro Gly Leu Pro Cys Pro Cys Thr Gly

5/27

	100	105	110
	Val Ser Asp Cys Ser Gly Gly Thr Asp Lys Lys Leu Arg Asn Cys Ser		
	115	120	125
	Arg Leu Ala Cys Leu Ala Gly Glu Leu Arg Cys Thr Leu Ser Asp Asp		
5	130	135	140
	Cys Ile Pro Leu Thr Trp Arg Cys Asp Gly His Pro Asp Cys Pro Asp		
	145	150	155
	Ser Ser Asp Glu Leu Gly Cys Gly Thr Asn Glu Ile Leu Pro Glu Gly		
	165	170	175
10	Asp Ala Thr Thr Met Gly Pro Pro Val Thr Leu Glu Ser Val Thr Ser		
	180	185	190
	Leu Arg Asn Ala Thr Thr Met Gly Pro Pro Val Thr Leu Glu Ser Val		
	195	200	205
	Pro Ser Val Gly Asn Ala Thr Ser Ser Ser Ala Gly Asp Gln Ser Gly		
15	210	215	220
	Ser Pro Thr Ala Tyr Gly Val Ile Ala Ala Ala Val Leu Ser Ala		
	225	230	235
	Ser Leu Val Thr Ala Thr Leu Leu Leu Leu Ser Trp Leu Arg Ala Gln		
	245	250	255
20	Glu Arg Leu Arg Pro Leu Gly Leu Leu Val Ala Met Lys Glu Ser Leu		
	260	265	270
	Leu Leu Ser Glu Gln Lys Thr Ser Leu Pro		
	275	280	
25	<210> 6		
	<211> 100		
	<212> PRT		
	<213> Homo sapiens		
30	<400> 6		
	Met Gly Asp Asp Gly Ser Ile Asp Tyr Thr Val His Glu Ala Trp Asn		
	1	5	10
			15

6/27

Glu Ala Thr Asn Val Tyr Leu Ile Val Ile Leu Val Ser Phe Gly Leu
 20 25 30
 Phe Met Tyr Ala Lys Arg Asn Lys Arg Arg Ile Met Arg Ile Phe Ser
 35 40 45
 5 Val Pro Pro Thr Glu Glu Thr Leu Ser Glu Pro Asn Phe Tyr Asp Thr
 50 55 60
 Ile Ser Lys Ile Arg Leu Arg Gln Gln Leu Glu Met Tyr Ser Ile Ser
 65 70 75 80
 Arg Lys Tyr Asp Tyr Gln Gln Pro Gln Asn Gln Ala Asp Ser Val Gln
 10 85 90 95
 Leu Ser Leu Glu
 100

 <210> 7
 15 <211> 225
 <212> PRT
 <213> Homo sapiens

 <400> 7
 20 Met Gly Thr Ala Asp Ser Asp Glu Met Ala Pro Glu Ala Pro Gln His
 1 5 10 15
 Thr His Ile Asp Val His Ile His Gln Glu Ser Ala Leu Ala Lys Leu
 20 25 30
 Leu Leu Thr Cys Cys Ser Ala Leu Arg Pro Arg Ala Thr Gln Ala Arg
 25 35 40 45
 Gly Ser Ser Arg Leu Leu Val Ala Ser Trp Val Met Gln Ile Val Leu
 50 55 60
 Gly Ile Leu Ser Ala Val Leu Gly Gly Phe Phe Tyr Ile Arg Asp Tyr
 65 70 75 80
 30 Thr Leu Leu Val Thr Ser Gly Ala Ala Ile Trp Thr Gly Ala Val Ala
 85 90 95
 Val Leu Ala Gly Ala Ala Ala Phe Ile Tyr Glu Lys Arg Gly Gly Thr

7/27

	100	105	110
	Tyr Trp Ala Leu Leu Arg Thr Leu Leu Ala Leu Ala Ala Phe Ser Thr		
	115	120	125
	Ala Ile Ala Ala Leu Lys Leu Trp Asn Glu Asp Phe Arg Tyr Gly Tyr		
5	130	135	140
	Ser Tyr Tyr Asn Ser Ala Cys Arg Ile Ser Ser Ser Ser Asp Trp Asn		
	145	150	155 160
	Thr Pro Ala Pro Thr Gln Ser Pro Glu Glu Val Arg Arg Leu His Leu		
	165	170	175
10	Cys Thr Ser Phe Met Asp Met Leu Lys Ala Leu Phe Arg Thr Leu Gln		
	180	185	190
	Ala Met Leu Leu Gly Val Trp Ile Leu Leu Leu Leu Ala Ser Leu Ala		
	195	200	205
	Pro Leu Trp Leu Tyr Cys Trp Arg Met Phe Pro Thr Lys Gly Val Ser		
15	210	215	220
	Pro		
	225		
	<210> 8		
20	<211> 387		
	<212> DNA		
	<213> Homo sapiens		
	<400> 8		
25	atgcatttct tattcagatt cattgttttc ttttatctgt ggggcctttt tactgctcag	60	
	agacaaaaga aagaggagag caccgaagaa gtgaaaatag aagttttgca tcgtccagaa	120	
	aactgctcta agacaagcaa gaagggagac ctactaaatg cccattatga cggctacctg	180	
	gctaaagacg gctcgaaatt ctactgcagc cggacacaaa atgaaggcca ccccaaattg	240	
	tttgttcttg gtgttgggca agtcataaaa ggcctagaca ttgctatgac agatatgtgc	300	
30	cctggagaaa agcgaaaagt agttataccc ccttcatttg catacggaaa ggaaggctat	360	
	gataaacctc tacttgcaaa gggaatt	387	

8/27

<210> 9

<211> 405

<212> DNA

<213> Homo sapiens

5

<400> 9

atggtgctgg	agagcgtggc	cogtatcgtg	aaggtgcagc	tccctgcata	tctgaagcgg	60
ctcccagtc	ctgaaagcat	taccgggttc	gctaggctca	cagtttcaga	atggcttcgg	120
ttattgcctt	tccttgggtg	actcgcaact	cttggctacc	ttgcagttcg	tccattcctc	180
ccgaagaaga	aacaacagaa	ggatagcttg	attaatctta	aaatacaaaa	ggaaaatccg	240
aaagtagtga	atgaaataaa	cattgaagat	ttgtgtctta	ctaaagcagc	ttattgtagg	300
tgttggcggt	ctaaaacgtt	tcctgcctgc	gatgggtcac	ataataaaca	caatgaattg	360
acaggagata	atgtgggtcc	actaatactg	aagaagaaag	aagta		405

15

<210> 10

<211> 237

<212> DNA

<213> Homo sapiens

20

<400> 10

atgccggtgg	ccgtgggtcc	ctacggacag	tcccagccaa	gctgcttcga	ccgtgtcaaa	60
atgggcttcg	tgatgggttg	cgccgtgggc	atggcgcccg	gggcgctctt	cggcaccttt	120
tcctgtctca	ggatcggaat	gcggggtcga	gagctgatgg	gcggcattgg	gaaaaccatg	180
atgcagagtg	gcggcacctt	tggcacattc	atggccattg	ggatgggcat	ccgatgc	237

25

<210> 11

<211> 432

<212> DNA

<213> Homo sapiens

30

<400> 11

atggcggtta	cgttcgccgc	cttctgctac	atgetggcgc	tgctgctcac	tgccgcgctc	60
------------	------------	------------	------------	------------	------------	----

9/27

	atctttcttcg ccatttggca cattatagca tttgatgagc tgaagactga ttacaagaat	120
	cctatagacc agtgtaatac cctgaatccc cttgtactcc cagagtacct catccacgct	180
	ttctttctgtg tcatgtttct ttgtgcagca gagggtgcta cactgggtct caatatgccc	240
	ctcttggcat atcatatttg gaggtatatg agtagaccag tgatgagtgg cccaggactc	300
5	tatgacccta caaccatcat gaatgcagat attctagcat attgtcagaa ggaaggatgg	360
	tgcaaattag ctttttatct tctagcattt ttttactacc tatatggcat gatctatgtt	420
	ttggtgagct ct	432
	<210> 12	
10	<211> 846	
	<212> DNA	
	<213> Homo sapiens	
	<400> 12	
15	atgagcggcg gttggatggc gcaggttgga gcgtggcgaa caggggctct gggcctggcg	60
	ctgctgctgc tgctcgccct cggactaggc ctggaggccg ccgcgagccc gctttccacc	120
	ccgacctctg cccaggccgc agggcccagc tcaggctcgt gccacccac caagttccag	180
	tgccgcacca gtggcttatg cgtgcccctc acctggcgct gcgacaggga cttggactgc	240
	agcgatggca gcgatgagga ggagtgcagg attgagccat gtaccagaa agggcaatgc	300
20	ccaccgcccc ctggcctccc ctgcccctgc accggcgcta gtgactgctc tgggggaact	360
	gacaagaaac tgcgcaactg cagccgcctg gcctgcctag caggcgagct ccgttgcacg	420
	ctgagcgatg actgcattcc actcacgtgg cgctgcgacg gccaccagaa ctgtcccac	480
	tccagcgacg agctcggctg tggaaccaat gagatcctcc cggaagggga tgccacaacc	540
	atggggcccc ctgtgacctt ggagagtgtc acctctctca ggaatgccac aaccatgggg	600
25	ccccctgtga ccctggagag tgtcccctct gtcggaatg ccacatcctc ctctgccgga	660
	gaccagtctg gaagcccaac tgcctatggg gttattgcag ctgctgcggt gctcagtga	720
	agcctggtca ccgccaccct cctccttttg tcctggtctc gagcccagga gcgcctccgc	780
	ccactggggt tactggtggc catgaaggag tccctgctgc tgtcagaaca gaagacctcg	840
	ctgccc	846
30	<210> 13	
	<211> 300	

10/27

<212> DNA

<213> Homo sapiens

<400> 13

5	atgggtgacg atggttctat tgattatact gttcacgaag cctggaatga agccaccaat	60
	gtttacttga tagttatcct tgtagcttc ggtctcttca tgtatgccaa aaggaacaaa	120
	aggagaatta tgaggatatt cagtgtgcc cctacagagg aaactttgtc agagcccaac	180
	ttttatgaca cgataagcaa gattcgttta agacaacaac tggaaatgta ttccatttca	240
	agaaagtacg actatcagca gccacaaaac caagctgaca gtgtgcaact ctcattggaa	300

10

<210> 14

<211> 675

<212> DNA

<213> Homo sapiens

15

<400> 14

20	atgggaacag ccgacagtga tgagatggcc ccggaggccc cacagcacac ccacatcgat	60
	gtgcacatcc accaggagtc tgccctggcc aagctcctgc tcacctgctg ctctgcgctg	120
	cggccccggg ccaccaggc caggggcagc agccggctgc tggtagcctc gtgggtgatg	180
	cagatcgtgc tggggatctt gagtgcagtc ctaggaggat ttttctacat ccgcgactac	240
	accctcctcg tcacctcggg agctgccatc tggacagggg ctgtggctgt gctggctgga	300
	gctgtgcct tcatttacga gaaacggggg ggtacatact gggccctgct gaggactctg	360
	ctagcgtgg cagctttctc cacagccatc gctgccctca aactttggaa tgaagatttc	420
	cgatatggct actcttatta caacagtgcc tgccgcatct ccagctcgag tgactggaac	480
25	actccagccc ccactcagag tccagaagaa gtcagaaggc tacacctatg tacctccttc	540
	atggacatgc tgaaggcctt gttcagaacc ctccaggcca tgctcttggg tgtctggatt	600
	ctgctgcttc tggcatctct ggccccctctg tggctgtact gctggagaat gttcccaacc	660
	aaagggtga gtccc	675

30

<210> 15

<211> 761

<212> DNA

11/27

<213> Homo sapiens

<400> 15

	gacatccacg gggcgcgagt gacacgcggg agggagagca gtgttctgct ggagccgatg	60
5	ccaaaaacc atg cat ttc tta ttc aga ttc att gtt ttc ttt tat ctg tgg	111
	Met His Phe Leu Phe Arg Phe Ile Val Phe Phe Tyr Leu Trp	
	1 5 10	
	ggc ctt ttt act gct cag aga caa aag aaa gag gag agc acc gaa gaa	159
	Gly Leu Phe Thr Ala Gln Arg Gln Lys Lys Glu Glu Ser Thr Glu Glu	
10	15 20 25 30	
	gtg aaa ata gaa gtt ttg cat cgt cca gaa aac tgc tct aag aca agc	207
	Val Lys Ile Glu Val Leu His Arg Pro Glu Asn Cys Ser Lys Thr Ser	
	35 40 45	
	aag aag gga gac cta cta aat gcc cat tat gac ggc tac ctg gct aaa	255
15	Lys Lys Gly Asp Leu Leu Asn Ala His Tyr Asp Gly Tyr Leu Ala Lys	
	50 55 60	
	gac ggc tcg aaa ttc tac tgc agc cgg aca caa aat gaa ggc cac ccc	303
	Asp Gly Ser Lys Phe Tyr Cys Ser Arg Thr Gln Asn Glu Gly His Pro	
	65 70 75	
20	aaa tgg ttt gtt ctt ggt gtt ggg caa gtc ata aaa ggc cta gac att	351
	Lys Trp Phe Val Leu Gly Val Gly Gln Val Ile Lys Gly Leu Asp Ile	
	80 85 90	
	gct atg aca gat atg tgc cct gga gaa aag cga aaa gta gtt ata ccc	399
	Ala Met Thr Asp Met Cys Pro Gly Glu Lys Arg Lys Val Val Ile Pro	
25	95 100 105 110	
	cct tca ttt gca tac gga aag gaa ggc tat gat aaa cct cta ctt gca	447
	Pro Ser Phe Ala Tyr Gly Lys Glu Gly Tyr Asp Lys Pro Leu Leu Ala	
	115 120 125	
	aag gga att tgaaaaagat gagaagccac gtgacaagtc atatcaggat gcag	500
30	Lys Gly Ile	
	ttttagaaga tattttttaag aagaatgacc atgatggtga tggcttcatt tctcccaagg	560

12/27

aatacaatgt ataccaacac gatgaactat agcatatttg tatttctact ttttttttta 620
 gctattttact gtactttatg tataaaacaa agtcactttt ctccaagttg tatttgctat 680
 ttttccccta tgagaagata ttttgatctc cccaatacat tgattttggt ataataaaat 740
 gtgaggctgt tttgcaaact t 761

5

<210> 16
 <211> 129
 <212> PRT
 <213> Homo sapiens

10

<400> 16

Met His Phe Leu Phe Arg Phe Ile Val Phe Phe Tyr Leu Trp

1 5 10

Gly Leu Phe Thr Ala Gln Arg Gln Lys Lys Glu Glu Ser Thr Glu Glu

15 15 20 25 30

Val Lys Ile Glu Val Leu His Arg Pro Glu Asn Cys Ser Lys Thr Ser

35 40 45

Lys Lys Gly Asp Leu Leu Asn Ala His Tyr Asp Gly Tyr Leu Ala Lys

50 55 60

20 Asp Gly Ser Lys Phe Tyr Cys Ser Arg Thr Gln Asn Glu Gly His Pro

65 70 75

Lys Trp Phe Val Leu Gly Val Gly Gln Val Ile Lys Gly Leu Asp Ile

80 85 90

Ala Met Thr Asp Met Cys Pro Gly Glu Lys Arg Lys Val Val Ile Pro

25 95 100 105 110

Pro Ser Phe Ala Tyr Gly Lys Glu Gly Tyr Asp Lys Pro Leu Leu Ala

115 120 125

Lys Gly Ile

30

<210> 17

<211> 701

<212> DNA

13/27

<213> Homo sapiens

<400> 17

```

cagagcggag ggggctcggg agaggagtgg acgccgctgg ccagg atg gtg ctg gag      57
5                                     Met Val Leu Glu
                                     1

agc gtg gcc cgt atc gtg aag gtg cag ctc cct gca tat ctg aag cgg      105
Ser Val Ala Arg Ile Val Lys Val Gln Leu Pro Ala Tyr Leu Lys Arg
      5              10              15              20

10 ctc cca gtc cct gaa agc att acc ggg ttc gct agg ctc aca gtt tca      153
Leu Pro Val Pro Glu Ser Ile Thr Gly Phe Ala Arg Leu Thr Val Ser
              25              30              35

gaa tgg ctt cgg tta ttg cct ttc ctt ggt gta ctc gca ctt ctt ggc      201
Glu Trp Leu Arg Leu Leu Pro Phe Leu Gly Val Leu Ala Leu Leu Gly
15              40              45              50

tac ctt gca gtt cgt cca ttc ctc ccg aag aag aaa caa cag aag gat      249
Tyr Leu Ala Val Arg Pro Phe Leu Pro Lys Lys Lys Gln Gln Lys Asp
              55              60              65

agc ttg att aat ctt aaa ata caa aag gaa aat ccg aaa gta gtg aat      297
20 Ser Leu Ile Asn Leu Lys Ile Gln Lys Glu Asn Pro Lys Val Val Asn
      70              75              80

gaa ata aac att gaa gat ttg tgt ctt act aaa gca gct tat tgt agg      345
Glu Ile Asn Ile Glu Asp Leu Cys Leu Thr Lys Ala Ala Tyr Cys Arg
      85              90              95              100

25 tgt tgg cgt tct aaa acg ttt cct gcc tgc gat ggt tca cat aat aaa      393
Cys Trp Arg Ser Lys Thr Phe Pro Ala Cys Asp Gly Ser His Asn Lys
              105              110              115

cac aat gaa ttg aca gga gat aat gtg ggt cca cta ata ctg aag aag      441
His Asn Glu Leu Thr Gly Asp Asn Val Gly Pro Leu Ile Leu Lys Lys
30              120              125              130

aaa gaa gta taataataat aacaatattt tctcattcctt tgtgtataga      490
Lys Glu Val

```

14/27

135

aaattttaaa atggtggtct taattattac tactggttga acaattattt cttccaattt 550
 atttttcttc tgcactactg tttgtatttg atcctttgtc tattcagtca ctttaattaga 610
 aattaaattg tcaagcctct tattctgact tcaaagaatt aatgtatctt ccaacaataa 670
 5 aatcacttct gattttaatc taggaaaacc t 701

<210> 18

<211> 135

<212> PRT

10 <213> Homo sapiens

<400> 18

Met Val Leu Glu

1

15 Ser Val Ala Arg Ile Val Lys Val Gln Leu Pro Ala Tyr Leu Lys Arg
 5 10 15 20
 Leu Pro Val Pro Glu Ser Ile Thr Gly Phe Ala Arg Leu Thr Val Ser
 25 30 35
 Glu Trp Leu Arg Leu Leu Pro Phe Leu Gly Val Leu Ala Leu Leu Gly
 20 40 45 50
 Tyr Leu Ala Val Arg Pro Phe Leu Pro Lys Lys Lys Gln Gln Lys Asp
 55 60 65
 Ser Leu Ile Asn Leu Lys Ile Gln Lys Glu Asn Pro Lys Val Val Asn
 70 75 80
 25 Glu Ile Asn Ile Glu Asp Leu Cys Leu Thr Lys Ala Ala Tyr Cys Arg
 85 90 95 100
 Cys Trp Arg Ser Lys Thr Phe Pro Ala Cys Asp Gly Ser His Asn Lys
 105 110 115
 His Asn Glu Leu Thr Gly Asp Asn Val Gly Pro Leu Ile Leu Lys Lys
 30 120 125 130
 Lys Glu Val

135

15/27

<210> 19
 <211> 393
 <212> DNA
 5 <213> Homo sapiens

<400> 19
 tcacgcgggcc gcgagcgccc tccccgtcgt tttccgtgag agacgtagag ctgagcgacc 60
 cagcccgcga gcgaggtgag atg ccg gtg gcc gtg ggt ccc tac gga cag tcc 113
 10 Met Pro Val Ala Val Gly Pro Tyr Gly Gln Ser
 1 5 10
 cag cca agc tgc ttc gac cgt gtc aaa atg gcc ttc gtg atg ggt tgc 161
 Gln Pro Ser Cys Phe Asp Arg Val Lys Met Gly Phe Val Met Gly Cys
 15 20 25
 15 gcc gtg ggc atg gcg gcc ggg gcg ctc ttc ggc acc ttt tcc tgt ctc 209
 Ala Val Gly Met Ala Ala Gly Ala Leu Phe Gly Thr Phe Ser Cys Leu
 30 35 40
 agg atc gga atg cgg ggt cga gag ctg atg gcc gcc att ggg aaa acc 257
 Arg Ile Gly Met Arg Gly Arg Glu Leu Met Gly Gly Ile Gly Lys Thr
 20 45 50 55
 atg atg cag agt gcc gcc acc ttt gcc aca ttc atg gcc att ggg atg 305
 Met Met Gln Ser Gly Gly Thr Phe Gly Thr Phe Met Ala Ile Gly Met
 60 65 70 75
 gcc atc cga tgc taaccatggt tgccaactac atctgtccct tcc 350
 25 Gly Ile Arg Cys
 gcc atc cga tgc taaccatggt tgccaactac atctgtccct tcccatcaat ccc 360
 Gly Ile Arg Cys
 agcccatgta ctaataaaaag aaagtctttg agt 393

30 <210> 20
 <211> 83
 <212> PRT

<213> Homo sapiens

Met Pro Val Ala Val Gly Pro Tyr Gly Gln Ser

[illegible]

<212> DNA

20 <213> Homo sapiens

<400> 21

	ctttctccgc tggcaacggc gccgctcccc gctcctcctc cccagcc atg gcg ttc	56
	Met Ala Phe	
25	1	
	acg ttc gcg gcc ttc tgc tac atg ctg gcg ctg ctg ctc act gcc gcg	104
	Thr Phe Ala Ala Phe Cys Tyr Met Leu Ala Leu Leu Leu Thr Ala Ala	
	5 10 15	
	ctc atc ttc ttc gcc att tgg cac att ata gca ttt gat gag ctg aag	152
30	Leu Ile Phe Phe Ala Ile Trp His Ile Ile Ala Phe Asp Glu Leu Lys	
	20 25 30 35	
	act gat tac aag aat cct ata gac cag tgt aat acc ctg aat ccc ctt	200

17/27

	Thr Asp Tyr Lys Asn Pro Ile Asp Gln Cys Asn Thr Leu Asn Pro Leu	
	40 45 50	
	gta ctc cca gag tac ctc atc cac gct ttc ttc tgt gtc atg ttt ctt	248
	Val Leu Pro Glu Tyr Leu Ile His Ala Phe Phe Cys Val Met Phe Leu	
5	55 60 65	
	tgt gca gca gag tgg ctt aca ctg ggt ctc aat atg ccc ctc ttg gca	296
	Cys Ala Ala Glu Trp Leu Thr Leu Gly Leu Asn Met Pro Leu Leu Ala	
	70 75 80	
	tat cat att tgg agg tat atg agt aga cca gtg atg agt ggc cca gga	344
10	Tyr His Ile Trp Arg Tyr Met Ser Arg Pro Val Met Ser Gly Pro Gly	
	85 90 95	
	ctc tat gac cct aca acc atc atg aat gca gat att cta gca tat tgt	392
	Leu Tyr Asp Pro Thr Thr Ile Met Asn Ala Asp Ile Leu Ala Tyr Cys	
	100 105 110 115	
15	cag aag gaa gga tgg tgc aaa tta gct ttt tat ctt cta gca ttt ttt	440
	Gln Lys Glu Gly Trp Cys Lys Leu Ala Phe Tyr Leu Leu Ala Phe Phe	
	120 125 130	
	tac tac cta tat ggc atg atc tat gtt ttg gtg agc tct tagaacaaca c	490
	Tyr Tyr Leu Tyr Gly Met Ile Tyr Val Leu Val Ser Ser	
20	135 140	
	acagaagaat tgggtccagtt aagtgcagtc aaaaagccac caaatgaagg gattctatcc	550
	agcaagatcc tgtccaagag tagcctgtgg aatctgatca gttactttaaa aaaatgactc	610
	cttattttttt aaatgtttcc acattttttgc ttgtggaaaag actgttttca tatgttatac	670
	tcagataaag attttaaatg gtattacgta taaattaata taaaatgatt acctctgggtg	730
25	ttgacagggtt tgaacttgca cttcttaagg aacagccata atcctctgaa tgatgcatta	790
	attactgact gtcctagtag attggaagct tttgtttata ggaacttgta gggctcattt	850
	tggttttcatt gaaacagtat ctaattataa attagctgta gatatcagggt gcttctgatg	910
	aagtgaaaat gtatatctga ctagtgggaa acttcatggg tttcctcatc tgtcatgtcg	970
	atgattatat atggatacat ttacaaaaat aaaaagcggg aattttccct tcgcttgaat	1030
30	att	1033

<210> 22

18/27

<211> 144

<212> PRT

<213> Homo sapiens

5 <400> 22

Met Ala Phe

1

Thr Phe Ala Ala Phe Cys Tyr Met Leu Ala Leu Leu Leu Thr Ala Ala

5 10 15

10 Leu Ile Phe Phe Ala Ile Trp His Ile Ile Ala Phe Asp Glu Leu Lys

20 25 30 35

Thr Asp Tyr Lys Asn Pro Ile Asp Gln Cys Asn Thr Leu Asn Pro Leu

40 45 50

Val Leu Pro Glu Tyr Leu Ile His Ala Phe Phe Cys Val Met Phe Leu

15 55 60 65

Cys Ala Ala Glu Trp Leu Thr Leu Gly Leu Asn Met Pro Leu Leu Ala

70 75 80

Tyr His Ile Trp Arg Tyr Met Ser Arg Pro Val Met Ser Gly Pro Gly

85 90 95

20 Leu Tyr Asp Pro Thr Thr Ile Met Asn Ala Asp Ile Leu Ala Tyr Cys

100 105 110 115

Gln Lys Glu Gly Trp Cys Lys Leu Ala Phe Tyr Leu Leu Ala Phe Phe

120 125 130

Tyr Tyr Leu Tyr Gly Met Ile Tyr Val Leu Val Ser Ser

25 135 140

<210> 23

<211> 1270

<212> DNA

30 <213> Homo sapiens

<400> 23

19/27

	ccccgccccca	accccgcgcg	tgcgcgtgcg	cagggataag	agagcgggtct	ggacagcgcg	60
	tggccgggcgc	cgctgtgggg	acagc	atg	agc	ggc ggt tgg atg gcg cag gtt	112
	Met Ser Gly Gly Trp Met Ala Gln Val						
	1		5				
5	gga gcg tgg cga aca ggg gct ctg ggc ctg gcg ctg ctg ctg ctg ctc						160
	Gly Ala Trp Arg Thr Gly Ala Leu Gly Leu Ala Leu Leu Leu Leu Leu						
	10	15	20	25			
	ggc ctc gga cta ggc ctg gag gcc gcc gcg agc ccg ctt tcc acc ccg						208
	Gly Leu Gly Leu Gly Leu Glu Ala Ala Ala Ser Pro Leu Ser Thr Pro						
10	30	35	40				
	acc tct gcc cag gcc gca ggc ccc agc tca ggc tcg tgc cca ccc acc						256
	Thr Ser Ala Gln Ala Ala Gly Pro Ser Ser Gly Ser Cys Pro Pro Thr						
	45	50	55				
	aag ttc cag tgc cgc acc agt ggc tta tgc gtg ccc ctc acc tgg cgc						304
15	Lys Phe Gln Cys Arg Thr Ser Gly Leu Cys Val Pro Leu Thr Trp Arg						
	60	65	70				
	tgc gac agg gac ttg gac tgc agc gat ggc agc gat gag gag gag tgc						352
	Cys Asp Arg Asp Leu Asp Cys Ser Asp Gly Ser Asp Glu Glu Glu Cys						
	75	80	85				
20	agg att gag cca tgt acc cag aaa ggg caa tgc cca ccg ccc cct ggc						400
	Arg Ile Glu Pro Cys Thr Gln Lys Gly Gln Cys Pro Pro Pro Pro Gly						
	90	95	100	105			
	ctc ccc tgc ccc tgc acc ggc gtc agt gac tgc tct ggg gga act gac						448
	Leu Pro Cys Pro Cys Thr Gly Val Ser Asp Cys Ser Gly Gly Thr Asp						
25	110	115	120				
	aag aaa ctg cgc aac tgc agc cgc ctg gcc tgc cta gca ggc gag ctc						496
	Lys Lys Leu Arg Asn Cys Ser Arg Leu Ala Cys Leu Ala Gly Glu Leu						
	125	130	135				
	cgt tgc acg ctg agc gat gac tgc att cca ctc acg tgg cgc tgc gac						544
30	Arg Cys Thr Leu Ser Asp Asp Cys Ile Pro Leu Thr Trp Arg Cys Asp						
	140	145	150				
	ggc cac cca gac tgt ccc gac tcc agc gac gag ctc ggc tgt gga acc						592

20/27

	Gly His Pro Asp Cys Pro Asp Ser Ser Asp Glu Leu Gly Cys Gly Thr	
	155 160 165	
	aat gag atc ctc ccg gaa ggg gat gcc aca acc atg ggg ccc cct gtg	640
	Asn Glu Ile Leu Pro Glu Gly Asp Ala Thr Thr Met Gly Pro Pro Val	
5	170 175 180 185	
	acc ctg gag agt gtc acc tct ctc agg aat gcc acc acc atg ggg ccc	688
	Thr Leu Glu Ser Val Thr Ser Leu Arg Asn Ala Thr Thr Met Gly Pro	
	190 195 200	
	cct gtg acc ctg gag agt gtc ccc tct gtc ggg aat gcc aca tcc tcc	736
10	Pro Val Thr Leu Glu Ser Val Pro Ser Val Gly Asn Ala Thr Ser Ser	
	205 210 215	
	tct gcc gga gac cag tct gga agc cca act gcc tat ggg gtt att gca	784
	Ser Ala Gly Asp Gln Ser Gly Ser Pro Thr Ala Tyr Gly Val Ile Ala	
	220 225 230	
15	gct gct gcg gtg ctc agt gca agc ctg gtc acc gcc acc ctc ctc ctt	832
	Ala Ala Ala Val Leu Ser Ala Ser Leu Val Thr Ala Thr Leu Leu Leu	
	235 240 245	
	ttg tcc tgg ctc cga gcc cag gag cgc ctc cgc cca ctg ggg tta ctg	880
	Leu Ser Trp Leu Arg Ala Gln Glu Arg Leu Arg Pro Leu Gly Leu Leu	
20	250 255 260 265	
	gtg gcc atg aag gag tcc ctg ctg ctg tca gaa cag aag acc tcg ctg	928
	Val Ala Met Lys Glu Ser Leu Leu Leu Ser Glu Gln Lys Thr Ser Leu	
	270 275 280	
	ccc tgaggacaag cacttgccac caccgtcact cagccctggg cgtagccgg	980
25	Pro	
	acaggaggag agcagtgatg cggatgggta cccgggcaca ccagccctca gagacctgag	1040
	ctcttctggc cacgtggaac ctggaacccg agctcctgca gaagtggccc tggagattga	1100
	gggtccctgg acaactcccta tggagatccg gggagctagg atggggaacc tgccacagcc	1160
30	agaactgagg ggctggcccc aggcagctcc cagggggtag aacggccctg tgcttaagac	1220
	actcctgctg ccccgctctga ggggtggcgat taaagttgct tcacatcctc	1270

```
<210> 24
<211> 282
<212> PRT
<213> Homo sapiens
```

<400> 24

[illegible]

	Pro Val Thr Leu Glu Ser Val Pro Ser Val Gly Asn Ala Thr Ser Ser	
	205	210 215
	Ser Ala Gly Asp Gln Ser Gly Ser Pro Thr Ala Tyr Gly Val Ile Ala	
	220	225 230
5	Ala Ala Ala Val Leu Ser Ala Ser Leu Val Thr Ala Thr Leu Leu Leu	
	235	240 245
	Leu Ser Trp Leu Arg Ala Gln Glu Arg Leu Arg Pro Leu Gly Leu Leu	
	250	255 260 265
	Val Ala Met Lys Glu Ser Leu Leu Leu Ser Glu Gln Lys Thr Ser Leu	
10	270	275 280
	Pro	
	<210> 25	
	<211> 836	
15	<212> DNA	
	<213> Homo sapiens	
	<400> 25	
	aaaaaaaaagg aaatgacgaa ggcagagggc gtccaggtcc gctcggtaac cgtttcccg	60
20	gcgcccggcc ccgactccgg ggtaaagagc cccggagcgg agcagcgctg gccgcgtgcc	120
	gcctccggag ccggcagccc ccatggctgg gggttatgga gtg atg ggt gac gat	175
	Met Gly Asp Asp	
	1	
	ggt tct att gat tat act gtt cac gaa gcc tgg aat gaa gcc acc aat	223
25	Gly Ser Ile Asp Tyr Thr Val His Glu Ala Trp Asn Glu Ala Thr Asn	
	5 10 15 20	
	gtt tac ttg ata gtt atc ctt gtt agc ttc ggt ctc ttc atg tat gcc	271
	Val Tyr Leu Ile Val Ile Leu Val Ser Phe Gly Leu Phe Met Tyr Ala	
	25 30 35	
30	aaa agg aac aaa agg aga att atg agg ata ttc agt gtg cca cct aca	319
	Lys Arg Asn Lys Arg Arg Ile Met Arg Ile Phe Ser Val Pro Pro Thr	
	40 45 50	

23/27

	gag gaa act ttg tca gag ccc aac ttt tat gac acg ata agc aag att	367
	Glu Glu Thr Leu Ser Glu Pro Asn Phe Tyr Asp Thr Ile Ser Lys Ile	
	55 60 65	
	cgt tta aga caa caa ctg gaa atg tat tcc att tca aga aag tac gac	415
5	Arg Leu Arg Gln Gln Leu Glu Met Tyr Ser Ile Ser Arg Lys Tyr Asp	
	70 75 80	
	tat cag cag cca caa aac caa gct gac agt gtg caa ctc tca ttg gaa	463
	Tyr Gln Gln Pro Gln Asn Gln Ala Asp Ser Val Gln Leu Ser Leu Glu	
	85 90 95 100	
10	tgaaacc tcagaaaaag agcaacagaa gtaattgttt caagctcctg attctttcta	520
	ctaaatcatg aacagcttta aaaacatttc tgtctgcata aaattatattt acttgtaact	580
	tttccccaat tgttctgtgc attgttttgc ctttttaaata tacatctcca agtggctcaa	640
	aaggccttga cacaggggaac ctgcacatat ccaggatatg tgtaaccagc gatggtgact	700
	tgaccttgcc aagacctgtg attccttcag gatacaatca gtgagaaata aaaacacatc	760
15	ttgggaagtg ggaatcctgg agtttatgcc atttgcaata ttaaaaaata aaaatgcaag	820
	ttattatttc aataat	836
	<210> 26	
	<211> 100	
20	<212> PRT	
	<213> Homo sapiens	
	<400> 26	
		Met Gly Asp Asp
25		1
	Gly Ser Ile Asp Tyr Thr Val His Glu Ala Trp Asn Glu Ala Thr Asn	
	5 10 15 20	
	Val Tyr Leu Ile Val Ile Leu Val Ser Phe Gly Leu Phe Met Tyr Ala	
	25 30 35	
30	Lys Arg Asn Lys Arg Arg Ile Met Arg Ile Phe Ser Val Pro Pro Thr	
	40 45 50	
	Glu Glu Thr Leu Ser Glu Pro Asn Phe Tyr Asp Thr Ile Ser Lys Ile	

24/27

	55	60	65	
	Arg Leu Arg Gln Gln Leu Glu Met Tyr Ser Ile Ser Arg Lys Tyr Asp			
	70	75	80	
	Tyr Gln Gln Pro Gln Asn Gln Ala Asp Ser Val Gln Leu Ser Leu Glu			
5	85	90	95	100
 <210> 27				
<211> 1022				
<212> DNA				
10	<213> Homo sapiens			
 <400> 27				
	agccctcccg ccgcccgcgc gcaggtcccg aggagcgag actgtgtccc tgaca atg			58
				Met
15				1
	gga aca gcc gac agt gat gag atg gcc ccg gag gcc cca cag cac acc			106
	Gly Thr Ala Asp Ser Asp Glu Met Ala Pro Glu Ala Pro Gln His Thr			
	5	10	15	
	cac atc gat gtg cac atc cac cag gag tct gcc ctg gcc aag ctc ctg			154
20	His Ile Asp Val His Ile His Gln Glu Ser Ala Leu Ala Lys Leu Leu			
	20	25	30	
	ctc acc tgc tgc tct gcg ctg cgg ccc cgg gcc acc cag gcc agg ggc			202
	Leu Thr Cys Cys Ser Ala Leu Arg Pro Arg Ala Thr Gln Ala Arg Gly			
	35	40	45	
25	agc agc cgg ctg ctg gtg gcc tcg tgg gtg atg cag atc gtg ctg ggg			250
	Ser Ser Arg Leu Leu Val Ala Ser Trp Val Met Gln Ile Val Leu Gly			
	50	55	60	65
	atc ttg agt gca gtc cta gga gga ttt ttc tac atc cgc gac tac acc			298
	Ile Leu Ser Ala Val Leu Gly Gly Phe Phe Tyr Ile Arg Asp Tyr Thr			
30	70	75	80	
	ctc ctc gtc acc tcg gga gct gcc atc tgg aca ggg gct gtg gct gtg			346
	Leu Leu Val Thr Ser Gly Ala Ala Ile Trp Thr Gly Ala Val Ala Val			

25/27

	85	90	95	
	ctg gct gga gct gct gcc ttc att tac gag aaa cgg ggt ggt aca tac			394
	Leu Ala Gly Ala Ala Ala Phe Ile Tyr Glu Lys Arg Gly Gly Thr Tyr			
	100	105	110	
5	tgg gcc ctg ctg agg act ctg cta gcg ctg gca gct ttc tcc aca gcc			442
	Trp Ala Leu Leu Arg Thr Leu Leu Ala Leu Ala Ala Phe Ser Thr Ala			
	115	120	125	
	atc gct gcc ctc aaa ctt tgg aat gaa gat ttc cga tat ggc tac tct			490
	Ile Ala Ala Leu Lys Leu Trp Asn Glu Asp Phe Arg Tyr Gly Tyr Ser			
10	130	135	140	145
	tat tac aac agt gcc tgc cgc atc tcc agc tcg agt gac tgg aac act			538
	Tyr Tyr Asn Ser Ala Cys Arg Ile Ser Ser Ser Ser Asp Trp Asn Thr			
	150	155	160	
	cca gcc ccc act cag agt cca gaa gaa gtc aga agg cta cac cta tgt			586
15	Pro Ala Pro Thr Gln Ser Pro Glu Glu Val Arg Arg Leu His Leu Cys			
	165	170	175	
	acc tcc ttc atg gac atg ctg aag gcc ttg ttc aga acc ctt cag gcc			634
	Thr Ser Phe Met Asp Met Leu Lys Ala Leu Phe Arg Thr Leu Gln Ala			
	180	185	190	
20	atg ctc ttg ggt gtc tgg att ctg ctg ctt ctg gca tct ctg gcc cct			682
	Met Leu Leu Gly Val Trp Ile Leu Leu Leu Leu Ala Ser Leu Ala Pro			
	195	200	205	
	ctg tgg ctg tac tgc tgg aga atg ttc cca acc aaa ggg gtg agt ccc			730
	Leu Trp Leu Tyr Cys Trp Arg Met Phe Pro Thr Lys Gly Val Ser Pro			
25	210	215	220	225
	taagaaaaga gaccagaagg aaatgttgga agtgagtgga atctagccat gcctctcctg			790
	attattagtg cctggtgctt ctgcaccggg cgtccctgca tctgactgct ggaagaagaa			850
	ccagactgag gaaaagaggc tcttcaacag cccagttat cctggcccca tgaccgtggc			910
	cacagccctg ctccagcagc acttgcccat tccttacacc ccttccccat cctgctccgc			970
30	ttcatgtccc ctccctgagta gtcattgtgat aataaactct catgttattg tt			1022

26/27

<211> 225

<212> PRT

<213> Homo sapiens

5 <400> 28

Met

1

Gly Thr Ala Asp Ser Asp Glu Met Ala Pro Glu Ala Pro Gln His Thr
 5 10 15
 10 His Ile Asp Val His Ile His Gln Glu Ser Ala Leu Ala Lys Leu Leu
 20 25 30
 Leu Thr Cys Cys Ser Ala Leu Arg Pro Arg Ala Thr Gln Ala Arg Gly
 35 40 45
 Ser Ser Arg Leu Leu Val Ala Ser Trp Val Met Gln Ile Val Leu Gly
 15 50 55 60 65
 Ile Leu Ser Ala Val Leu Gly Gly Phe Phe Tyr Ile Arg Asp Tyr Thr
 70 75 80
 Leu Leu Val Thr Ser Gly Ala Ala Ile Trp Thr Gly Ala Val Ala Val
 85 90 95
 20 Leu Ala Gly Ala Ala Ala Phe Ile Tyr Glu Lys Arg Gly Gly Thr Tyr
 100 105 110
 Trp Ala Leu Leu Arg Thr Leu Leu Ala Leu Ala Ala Phe Ser Thr Ala
 115 120 125
 Ile Ala Ala Leu Lys Leu Trp Asn Glu Asp Phe Arg Tyr Gly Tyr Ser
 25 130 135 140 145
 Tyr Tyr Asn Ser Ala Cys Arg Ile Ser Ser Ser Ser Asp Trp Asn Thr
 150 155 160
 Pro Ala Pro Thr Gln Ser Pro Glu Glu Val Arg Arg Leu His Leu Cys
 165 170 175
 30 Thr Ser Phe Met Asp Met Leu Lys Ala Leu Phe Arg Thr Leu Gln Ala
 180 185 190
 Met Leu Leu Gly Val Trp Ile Leu Leu Leu Leu Ala Ser Leu Ala Pro

27/27

	195					200						205				
Leu	Trp	Leu	Tyr	Cys	Trp	Arg	Met	Phe	Pro	Thr	Lys	Gly	Val	Ser	Pro	
210						215					220					225



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, 5/10, C07K 14/705	A3	(11) International Publication Number: WO 99/43802 (43) International Publication Date: 2 September 1999 (02.09.99)
(21) International Application Number: PCT/JP99/00875 (22) International Filing Date: 25 February 1999 (25.02.99) (30) Priority Data: 10/46607 27 February 1998 (27.02.98) JP (71) Applicants (for all designated States except US): SAGAMI CHEMICAL RESEARCH CENTER [JP/JP]; 4-1, Nishi-Ohnuma 4-chome, Sagamihara-shi, Kanagawa 229-0012 (JP). PROTEGENE INC. [JP/JP]; 2-20-3, Naka-cho, Meguro-ku, Tokyo 153-0065 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): KATO, Seishi [JP/JP]; 3-46-50, Wakamatsu, Sagamihara-shi, Kanagawa 229-0014 (JP). SEKINE, Shingo [JP/JP]; Remonzu 101, 2-8-15, Atago, Ageo-shi, Saitama 362-0034 (JP). KIMURA, Tomoko [JP/JP]; 302, 4-1-28, Nishiikuta, Tama-ku, Kawasaki-shi, Kanagawa 214-0037 (JP). NAKAMURA, Nobuko [JP/JP]; 2-20-6, Kozonominami, Ayase-shi, Kanagawa 252-1122 (JP). (74) Agents: AOYAMA, Tamotsu et al.; Aoyama & Partners, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka 540-0001 (JP).		(81) Designated States: AU, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 29 December 1999 (29.12.99)
(54) Title: HUMAN PROTEINS HAVING TRANSMEMBRANE DOMAINS AND DNAs ENCODING THESE PROTEINS (57) Abstract Human proteins having transmembrane domains, cDNAs coding for these proteins, and expression vectors of said cDNAs as well as eucaryotic cells expressing said cDNAs. Said proteins and eucaryotic cells having said proteins on the membrane surface can be provided by expression of cDNAs coding for human proteins having transmembrane domains and of recombinants of these human cDNAs.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

Intern. Appl. Application No
PCT/JP 99/00875

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/12 C12N5/10 C07K14/705

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	<p>WO 98 54963 A (FERRIE ANN M; HUMAN GENOME SCIENCES INC ; GREENE JOHN M (US); YOUNG) 10 December 1998 see abstract sequence of gene 23 / clone HSQE084 (Human ortholog of mouse FKBP23): 100% identical to seq. ID 1 (aa 1-120). see abstract</p> <p style="text-align: center;">--- -/-</p>	1-6

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

15 July 1999

Date of mailing of the international search report

03. 11. 1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Galli, I

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/JP 99/00875

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE SEQUENCES EMBL, Heidelberg, FRG Accession No. AF040252, 3 February 1998 NAKAMURA T. ET AL.: "Mouse FK-506 binding protein (FKBP23)" XP002109279 amino acids 1-138 are 85.8% identical to seq. ID 1. see abstract & NAKAMURA T. ET AL.: "Molecular cloning, characterization and chromosomal localization of FKBP23, a novel FK506-binding protein with Ca²⁺-binding ability." GENOMICS, vol. 54, no. 1, November 1998, pages 89-98,</p>	1-6
X	<p>--- DATABASE GENBANK Accession No. AA457921, 6 June 1997 MARRA M. ET AL.: "EST; mouse cDNA clone IMAGE:849512 similar to FK506-binding protein precursor" XP002109280 the sequence encodes amino acids 24-100 of seq. ID 1 (93.9% identity) see abstract</p>	1-6
A	<p>--- WO 95 34641 A (SCRIPPS RESEARCH INST) 21 December 1995 see abstract see abstract; examples 1-4 see abstract; claims 1-47; examples 1-4</p>	1-6
A	<p>--- WO 93 23548 A (VERTEX PHARMA) 25 November 1993 see abstract; claims 1-47; examples 1-4</p>	1-6
A	<p>--- WO 95 00633 A (CHILDRENS MEDICAL CENTER ;UNIV LELAND STANFORD JUNIOR (US)) 5 January 1995 see abstract; claims 1-47; examples 1-4 -----</p>	1-6

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP 99/00875

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see FURTHER INFORMATION sheet, subject 1.

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/JP 99/00875

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: (1-6) - partial

Proteins containing the amino acid sequece of seq. ID 1.

Corresponding cDNAs (seq. IDs 8,15), vectors, transformed eucaryotic cells.

2. Claims: (1-6) - partial

Idem as subject matter 1 but limited to seq. IDs 2,9,17.

3. Claims: (1-6) - partial

Idem as subject matter 1 but limited to seq. IDs 3,10,19.

4. Claims: (1-6) - partial

Idem as subject matter 1 but limited to seq. IDs 4,11,21.

5. Claims: (1-6) - partial

Idem as subject matter 1 but limited to seq. IDs 5,12,23.

6. Claims: (1-6) - partial

Idem as subject matter 1 but limited to seq. IDs 6,13,25.

7. Claims: (1-6) - partial

Idem as subject matter 1 but limited to seq. IDs 7,14,27.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JP 99/00875

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9854963 A	10-12-1998	AU 7812098 A	21-12-1998
		AU 7811998 A	21-12-1998
		AU 6552198 A	29-09-1998
		WO 9840483 A	17-09-1998
WO 9534641 A	21-12-1995	AU 2773595 A	05-01-1996
WO 9323548 A	25-11-1993	AU 4388893 A	13-12-1993
WO 9500633 A	05-01-1995	US 5486599 A	23-01-1996
		AU 7112994 A	17-01-1995
		EP 0705332 A	10-04-1996